

RESEARCH ARTICLE

Comparison of “Live High-Train Low” in Normobaric versus Hypobaric Hypoxia

Jonas J. Saugy^{1,2}, Laurent Schmitt³, Roberto Cejuela⁴, Raphael Faiss^{1,2}, Anna Hauser^{1,5}, Jon P. Wehrlin⁵, Benjamin Rudaz¹, Audric Delessert¹, Neil Robinson⁶, Grégoire P. Millet^{1,2*}

1. ISSUL, Institute of Sport Sciences, Faculty of Biology and Medicine, University of Lausanne, Lausanne, Switzerland, 2. Department of Physiology, Faculty of Biology and Medicine, University of Lausanne, Lausanne, Switzerland, 3. National School of Mountain Sports/National Ski-Nordic Centre, Prémamanon, France, 4. Departmental Section of Physical Education and Sports, University of Alicante, Alicante, Spain, 5. Swiss Federal Institute of Sport, Magglingen, Switzerland, 6. Swiss Laboratory for Doping Analyses, University of Lausanne, Lausanne, Switzerland

*gregoire.millet@unil.ch



CrossMark
click for updates

OPEN ACCESS

Citation: Saugy JJ, Schmitt L, Cejuela R, Faiss R, Hauser A, et al. (2014) Comparison of “Live High-Train Low” in Normobaric versus Hypobaric Hypoxia. PLoS ONE 9(12): e114418. doi:10.1371/journal.pone.0114418

Editor: James West, Vanderbilt University Medical Center, United States of America

Received: September 6, 2014

Accepted: November 10, 2014

Published: December 17, 2014

Copyright: © 2014 Saugy et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability: The authors confirm that all data underlying the findings are fully available without restriction. All relevant data are within the paper.

Funding: This study was funded by grants from the Ministère des Sports, de la Jeunesse, de l'Éducation Populaire et de la Vie Associative (MSJEPVA)/Institut National du Sport, de l'Expertise et de la Performance (INSEP) and from Bundes Amt für Sport (BASPO; Switzerland).

Competing Interests: The authors have declared that no competing interests exist.

Abstract

We investigated the changes in both performance and selected physiological parameters following a Live High-Train Low (LHTL) altitude camp in either normobaric hypoxia (NH) or hypobaric hypoxia (HH) replicating current “real” practices of endurance athletes. Well-trained triathletes were split into two groups (NH, n=14 and HH, n=13) and completed an 18-d LHTL camp during which they trained at 1100–1200 m and resided at an altitude of 2250 m ($P_iO_2 = 121.7 \pm 1.2$ vs. 121.4 ± 0.9 mmHg) under either NH (hypoxic chamber; $F_iO_2 = 15.8 \pm 0.8\%$) or HH (real altitude; barometric pressure 580 ± 23 mmHg) conditions. Oxygen saturations (S_pO_2) were recorded continuously daily overnight. P_iO_2 and training loads were matched daily. Before (Pre-) and 1 day after (Post-) LHTL, blood samples, VO_{2max} , and total haemoglobin mass (Hb_{mass}) were measured. A 3-km running test was performed near sea level twice before, and 1, 7, and 21 days following LHTL. During LHTL, hypoxic exposure was lower for the NH group than for the HH group (220 vs. 300 h; $P < 0.001$). Night S_pO_2 was higher (92.1 ± 0.3 vs. $90.9 \pm 0.3\%$, $P < 0.001$), and breathing frequency was lower in the NH group compared with the HH group (13.9 ± 2.1 vs. 15.5 ± 1.5 breath.min⁻¹, $P < 0.05$). Immediately following LHTL, similar increases in VO_{2max} (6.1 ± 6.8 vs. $5.2 \pm 4.8\%$) and Hb_{mass} (2.6 ± 1.9 vs. $3.4 \pm 2.1\%$) were observed in NH and HH groups, respectively, while 3-km performance was not improved. However, 21 days following the LHTL intervention, 3-km run time was significantly faster in the HH ($3.3 \pm 3.6\%$; $P < 0.05$) versus the NH ($1.2 \pm 2.9\%$; ns) group. In conclusion, the greater degree of race performance enhancement by day 21 after an 18-d LHTL camp in the HH group was likely induced by a larger hypoxic dose. However, one cannot rule out other factors

including differences in sleeping desaturations and breathing patterns, thus suggesting higher hypoxic stimuli in the HH group.

Introduction

Live High - Train Low (LHTL) training camps are commonly used by athletes under either normobaric hypoxia (NH) [1–5] or hypobaric hypoxia (HH) [6–10] conditions. These two types of hypoxia are obtained by the combination of a lowered value of barometric pressure (PB) and/or a reduced inspired fraction of oxygen ($F_{I}O_2$) (NH: $F_{I}O_2 < 20.9\%$; PB=760 mmHg vs. HH: $F_{I}O_2 = 20.9\%$; PB<760 mmHg) resulting in an inspired partial pressure of oxygen ($P_{I}O_2$) less than 150 mmHg. NH and HH were, until recently, thought to be interchangeable since $P_{I}O_2$ was assumed as the only factor influencing the physiological responses to hypoxia [11]. This “*equivalent air altitude model*” [12] has now been criticized and a growing body of literature has reported physiological differences between acute exposures to NH and HH [13, 14]. Specifically, acute mountain sickness (AMS) symptoms are less severe under NH than HH conditions [15]. Pre-acclimatisation at real altitudes (HH) resulted in a significant decrease in the severity of AMS under HH conditions [16], which was not the case in individuals subjected to pre-acclimatisation under NH conditions [16]. Furthermore, according to Fulco et al., 2011, NH and HH could not “*be used interchangeably*” and do not exhibit the same levels of effectiveness relative to pre-acclimatisation strategies for the prevention of AMS (e.g., significant decrease in the severity of AMS under HH conditions following pre-acclimatization under HH but not NH) and for the improvement of exercise performance at higher altitudes [16]. In addition, minute ventilation was lower under HH than NH conditions with the combination of lower tidal volumes and higher respiratory frequencies [17]. Interestingly, oxidative stress markers were also elevated when individuals were continuously exposed to HH conditions compared to NH for 24 h, whereas nitric oxide (NO) in exhaled air and plasma was lower under HH *versus* NH [18]. Moreover, exhaled NO and NO end-products (NO_x) decreased in HH but remained stable in NH [18]. While the afore-mentioned studies support our recent suggestion that “*HH is a more severe environmental condition*” [14], this would also make the assumption that larger physiological adaptations would occur after prolonged hypoxic exposure under HH compared to NH conditions realistic. This still needs to be demonstrated in research.

Of interest is that the training practices of athletes are dependent upon hypoxic conditions, which reflect protocols described in the literature. It has indeed been reported that: 1) daily hypoxic exposure is shorter in NH (i.e. 8–12 h.d⁻¹; [4, 5]) compared to HH (i.e. 16–18 h.d⁻¹ in HH; [7, 19]) during LHTL protocols; 2) total hypoxic dose is reduced accordingly in NH (i.e. ~150–300 h; [1, 4, 5, 20, 21]) compared to HH (i.e. 300–600 h; [9, 22–25]); 3) total camp duration varies

between NH (11 to 23 days [2, 3, 26]; and HH (13 and 28 days; [9, 22–25]). Moreover, the mean performance improvements (*i.e.* power output increases) following LHTL were lower when the intervention was completed under NH compared to HH conditions (0.6% vs. 4.0%) [27]. Finally, most of the LHTL studies conducted under NH conditions did not elicit any performance improvement, although some induced positive erythropoietic responses [1, 28], and most of the studies reporting both performance and erythropoietic enhancements were performed under HH conditions [19, 29]. However, these findings of the differences in the physiological responses to NH vs. HH, which suggest larger adaptations in the HH condition, are based only on short-term hypoxic exposure. To the best of our knowledge, no study to date has directly compared altitude-induced adaptations (*i.e.* haematological, peripheral oxygen saturation, etc.) and performance changes following LHTL training camps under both NH and HH conditions. This is an important issue as the development of NH facilities (*e.g.*, nitrogen houses; hypoxic rooms, etc.) worldwide is increasing, and since coaches and athletes often consider NH and HH conditions to provide the same hypoxic stimulus. The issue of whether larger additional benefits occur by LHTL using HH conditions rather than NH conditions has never been directly investigated. Therefore, the present study aimed to compare the physiological responses and the performance gains in trained triathletes during and after LHTL camps with matched P_iO_2 in NH versus HH conditions. Of importance is that we replicated common or 'real' altitude training practices of endurance athletes (*e.g.*, daily exposure, total hypoxic doses under NH and HH conditions, respectively). We hypothesised that a LHTL intervention conducted under HH compared to NH should be advantageous for both physiological adaptation and performance increases.

Methods

Experimental Design

Our experimental design (Fig. 1) consisted of a 33-week period divided into the following four phases: 1) 24 weeks (January to May) were completed at sea level where training loads were quantified; followed by 2) a 3-wk lead-in period at sea level during which all training sessions were supervised and loads were quantified; 3) an 18-d LHTL training camp under either NH or HH conditions; and 4) a 3-wk post-altitude period at sea level during which all training sessions were once again supervised and loads were quantified.

Two groups (NH, $n=14$ and HH, $n=13$) were matched based on the VO_2 max values prior to the training camp and completed an 18-d LHTL camp during which all athletes trained at 1100–1200 m and resided at an altitude of 2250 m ($P_iO_2=121.7 \pm 1.2$ vs. 121.4 ± 0.9 mmHg) under either NH (hypoxic house; exposure 12.2 ± 0.3 h.d⁻¹; F_iO_2 $15.8 \pm 0.8\%$, Prémanson, France) or HH conditions (real altitude; 16.8 ± 3.1 h.d⁻¹; barometric pressure 580 ± 23 mmHg, Fiescheralp, Switzerland). Normobaric hypoxia was obtained by extracting oxygen




Fig. 1. Overview of the study design separated by weeks (W.) and in order of the testing altitude, including the six months before the lead-in period where the training loads were assessed, the lead-in, the LHTL camp of 18 days, and the lead-out period. Testing included 3-km test = the 3-km running tests on the track near sea level; Pre-, Post-, Post-7 and Post-21 = testing sessions; LHTL = Live High-Train Low training camp for normobaric hypoxia (NH) and hypobaric hypoxia (HH), where athletes were exposed to both higher altitude for living and sleeping and lower altitude for training.

doi:10.1371/journal.pone.0114418.g001

from ambient air (OBS, Husøysund, Norway). Calculations of $F_{I}O_2$ values corresponding to the required altitude took into account the altitude of Prémamon. As gas compositions were continuously monitored, O_2 fractions were permanently adjusted during sessions in order to maintain stability. Moreover, it was determined that opening the door several times for periods of a few seconds did not change $F_{I}O_2$ values. For safety reasons, O_2 and CO_2 compositions were monitored. Each room was connected to a central monitoring station under the control of an independent investigator.

The NH group was split into smaller groups and used hypoxic chambers, whereas the HH athletes went into the valley twice daily via cable car to perform the training.

Before (Pre-) and 24 h after (Post-) the LHTL period, several physiological tests were performed on both groups in the same location (Prémamon, France, 1150 m). These sessions were conducted in a well-ventilated laboratory (temperature $22 \pm 1^\circ C$) in the same order and at the same time of the day in the Pre- and Post- testing condition. [Measurements](#) included blood samples, anthropometric measurements, maximal incremental tests on a cycle ergometer (VO_{2max}), and total haemoglobin mass (Hb_{mass}) assessments. Subjects performed five 3-km running tests at the following times: prior to lead-in, before LHTL, after LHTL, seven days after LHTL (Post-7), and twenty one days following LHTL (Post-21). All 3-km running tests were performed near sea level (100–390 m).

Subjects

Twenty-seven well-trained male triathletes living at or near sea level (age 23 ± 4 years, body height 179 ± 5 cm, body weight 71 ± 7 kg, fat mass $10.1 \pm 1.6\%$, and VO_{2max} 66.9 ± 8.4 mL·kg⁻¹·min⁻¹) participated in this study. Three subjects were excluded following the lead-in period due to insufficient training loads and fitness. All subjects were non-smokers who had not been acclimatised or recently

exposed to testing altitudes. Volunteers provided their written, voluntary, informed consent before participation. The experiment was approved by a Medical Ethics Committee (Commission Cantonale Valaisanne d’Ethique Médicale, CCVEM; Agreement 051/09) and performed in accordance with the Declaration of Helsinki.

Measurements

Training content and training loads

Two experienced certified coaches supervised and advised the athletes during all training sessions of the lead-in period and matched training content and loads for both groups during the LHTL period. Additionally, each subject’s daily training loads were quantified individually by both subjective and objective means to evaluate their effects on physiological adaptation and each subject’s subsequent performance. Training load quantification was performed using the ‘Objective Load Scale’ (ECOs) [30]. The training loads for the triathletes included in our study were similar to those described in other studies involving endurance athletes. The Objective Load Scale allowed for the quantification of all training loads in each sport of the triathlon (swim, bike, run, and transitions). The daily and weekly training loads (ECOs) of each subject were quantified based on each subject’s physical characteristics and training program intensity. Volume was quantified by time and allowed for better comparisons of different performance levels and conditions (*e.g.*, ground surface, environmental temperature) [30].

3-km run test

The 3-km running tests were completed on a 400 m outdoor synthetic track near sea level. To avoid any group or pacing influences, starts were given in time-trial form (*e.g.*, 30 s between each start and randomisation of the order in which each athlete competed).

Incremental cycling test

VO_{2max} was tested before and after LHTL using subjects’ own bicycles, which were linked to a computerised ergometer system (Cyclus 2, RBM elektronik-automation GmbH, Leipzig, Germany). The exercise protocol began with a warm-up period of 5 min at a workload of 90 W. The workload was subsequently increased by $30 \text{ W}\cdot\text{min}^{-1}$ until voluntary exhaustion. During the final minutes of the test, subjects were strongly encouraged to perform until they reached maximal exhaustion and had achieved VO_{2max} based on the standard criteria for all tests. Each subject wore a mouthpiece and nose clip for breath collection. O_2 and CO_2 levels in expired gas were continuously measured and monitored as breath-by-breath values (Ultima Cardio 2 gas exchange analysis system, MGC Diagnostics with Breezesuite software, Saint Paul, MN, USA). Both the gas analyser and the flowmeter of the gas analyser were calibrated prior to each test. The highest 30 s

average value served as the VO_{2max} . Maximal heart rates (HR_{max}) and the lowest S_pO_2 values were each recorded during the same time period. The maximal power output (P_{max}) was the load of the last stage completed.

Anthropometrics values

Athletes' body weights and heights were measured in the morning before breakfast.

Haemoglobin mass

Hb_{mass} was measured in duplicate by using a slightly modified version [31] of the optimised carbon monoxide (CO)-rebreathing method described by Schmidt and Prommer [32]. The subjects inhaled a bolus of 100 mL of pure CO (Multigas SA, Domdidier, Switzerland), followed by 3.5 L of oxygen. Each Hb_{mass} measurement was performed in duplicate on two consecutive days (12- to 24-h time lag between measurements). Across all time points, the mean error for duplicate Hb_{mass} measurement was 2.1% in our mobile laboratory. Hb_{mass} data are expressed as the mean values of the duplicate measurements.

Values of red cell volume (RCV), blood volume (BV), and plasma volume (PV) were estimated using the following formulas: $RCV = Hb_{mass}/MCHC \times 100$, $BV = RCV \times (100/Hct)$ and $PV = BV - RCV$, where MCHC is the mean corpuscular haemoglobin concentration, and Hct is the haematocrit corrected to whole-body haematocrit by the cell factor of 0.91. For the calculation of RCV, BV, and PV, venous haemoglobin concentrations [Hb] and venous Hct were used. Blood gas analyses were conducted using an ABL 800flex (Radiometer A/S, Copenhagen, Denmark).

Blood samples

Antecubital vein blood samples (4.9 mL EDTA tube, Sarstedt, Nümbrecht, Germany) were taken during three time periods including either before breakfast or immediately after waking up, before LHTL, and after LHTL. Blood was subsequently analysed via fluorescent flow cytometry and hydrodynamic focusing (XT-2000i, Sysmex Europe, Norderstedt, Germany), and the following primary haematological parameters were quantified: red blood cells (RBC), haemoglobin (Hb), haematocrit (Hct), mean cell volume (MCV), mean cell haemoglobin (MCH), mean cell haemoglobin concentration (MCHC), reticulocyte percentage (RET%), absolute number of reticulocytes (RET#), and immature reticulocyte fraction (IRF). The Sysmex XT-2000i underwent regular internal quality control procedures as required by the standards of laboratory medicine. During the period of our study, the coefficient of variations (CV), which was determined using internal quality controls, was far below 1.5% for Hb and 1.5% for RET% (within CV limits accepted by the manufacturer of the instrument). Plasma EPO was quantified using a standard procedure with an ELISA kit (Stemcell

Technologies, Grenoble, France). Plasma EPO concentrations below the limit of quantification (1.6 mU/mL) were excluded from the analyses. CVs determined by three internal quality controls (levels: low, medium and high) were below 15% in our WADA (World Anti-Doping Agency) accredited laboratory [33]. All plasma samples were analysed in duplicate, where the mean values of the duplicate were used for this study. Additionally, baseline ferritin was quantified using standard laboratory procedures (Dimension EXL, Siemens Healthcare Diagnostics SA, Zürich, Switzerland) to evaluate the subject's iron stores. It is important to note that all athletes were tested for doping by the accredited laboratory according to the standards of the biological passport. This was done to avoid performance enhancement via doping.

Questionnaires

Subjects completed three different questionnaires on a daily basis immediately after waking up (hypoxic rooms for NH and normal rooms for HH, but all were hypoxic) and during three phases before, during, and after the LHTL training camp. The three questionnaires were as follows: 1) *The Lake Louise score questionnaire*, 2) *The Daily Analysis of Life Demands for Athletes (DALDA)*, and 3) *Sleep assessment questionnaire*.

The Lake Louise score questionnaire is scoring system developed by the 1991 International Hypoxia Symposium consensus committee, which met at Lake Louise in Alberta, Canada. It is widely used today to assess the severity of AMS. The DALDA is a self-reported sport-specific tool describing the stress sources and characteristics of each person, which allows for the differentiation of the individuality of stress responses. This questionnaire is divided into two parts; Part A describes the general stress sources that occur in everyday life for an athlete (diet, home life, school, work, friends, training, climate, sleep, recreation, and health); and Part B determines which symptoms of any existence in stress reactions of the athlete. The sleep assessment questionnaire was the Groningen Sleep Quality Scale (GSQS), which was used to evaluate for high altitude sleep (HAS) disturbances. It consists of a sleep quality score (GSQSS) and two visual analog scales (VAS), which yield a score between 0 and 10 for sleep quality and waking state.

Sleep assessment

S_pO_2 and HR were recorded each evening at 0.25 Hz with a wrist oximeter connected to a finger sensor (Wristox 3150 with 8000SM-WO Sensor, Nonin, Plymouth, MN). Subjects wore an instrumented t-shirt (model SEW, CSEM, Neuchâtel, Switzerland) each night (including the 2 nights before and the 2 nights after LHTL), a device made of comfortable fabric that was used to measure breathing frequency via an elastic sensor included in the textile, as well as each subject's sleeping position via accelerometers.

Data Analysis and Statistics

Data are reported as the means and standard deviations. Data were tested for equality of variance (Fisher-Snedecor *F-test*) and for normality (Shapiro-Wilk test). When both conditions were met, a two-way ANOVA was performed for repeated measures for each condition (NH and HH) to determine time effects for variables measured on several occasions during the camps with pairwise multiple comparison procedures (Holm-Sidak method). Differences between results obtained before and after LHTL for both the NH and HH groups were subsequently also compared using a two-way ANOVA. Differences in percentage changes between the groups were tested with a Wilcoxon signed rank sum test. When either equality of variance or normality were not satisfied, variables were analysed for each condition using a Friedman test for repeated measures to determine time effects using pairwise multiple comparison procedures (Bonferroni test). In this case, differences between the NH and HH groups at baseline (Pre-) were tested using a Mann-Whitney rank sum test. The correlation between values of Hb_{mass} initial (in g) or pre-to-post change (in %) and VO_{2max} (mL.kg⁻¹.min⁻¹) as well as correlations between all haematological and physiological parameters were calculated via the Pearson product moment correlation. Null hypotheses were rejected at $P < 0.05$. All analyses were completed using Sigmaplot 11.0 software (Systat Software, San Jose, CA).

Results

Training loads

No differences were found in daily or average training loads between the two groups during the 18-d LHTL camp (232 ± 159 vs. 217 ± 129 ECOs for the NH and HH groups, respectively; [Fig. 2A](#)). Additionally, no differences were found in weekly training loads monitored during the 6 months prior to the study (1161 ± 130 vs. 1208 ± 168 ECOs per week for the NH and HH groups, respectively; [Fig. 2B](#)), nor were any differences noted during the lead-in period or the post-hypoxic period.

3-km performance test

Compared to Pre-, 3-km performance remained unchanged at Post- and Post-7 in both groups. Whereas run performance did not improve significantly in the NH group (630.1 ± 64.8 vs. 621.8 ± 54.8 s, $P > 0.05$) from Pre- to Post-21, faster 3-km run times occurred in the HH group (611.1 ± 48.5 vs. 588.3 ± 32.2 s; $-3.3 \pm 3.6\%$, $P < 0.05$, [Fig. 3](#)). No differences were found between groups during the lead-in period, before LHTL, after LHTL, or 7 days post-LHTL. In addition, no differences were found between Lead-in and Pre- for both groups (626.3 ± 63.8 vs. 630.1 ± 64.8 s and 602.4 ± 44.3 vs. 611.1 ± 48.5 s, for NH and HH at Lead-in vs. Pre-, respectively).




Fig. 2. A. Daily objective training (D01-D18: day 01 to day 18) loads during the Live High-Train Low (LHTL) camp for normobaric hypoxia (NH) and hypobaric hypoxia (HH) groups. B. Weekly objective training loads during the six months before the intervention for NH and HH groups.

doi:10.1371/journal.pone.0114418.g002




Fig. 3. Relative change in 3-km run time from Pre- to Post-, Post-7, and Post-21 as determined on a running track near sea level for the normobaric hypoxia (NH) and hypobaric hypoxia (HH) groups (in %). Data are mean \pm standard error * $P < 0.05$ for differences with Pre- and # $P < 0.05$ for differences between groups.

doi:10.1371/journal.pone.0114418.g003

Maximal test on cycle ergometer

These results are presented in [Table 1](#). Both groups increased their maximal oxygen uptake values and power output values by the same amount immediately after the LHTL training camp period ($+6.1 \pm 6.8$ vs. $+5.2 \pm 4.8\%$ VO_{2max} and $+9.6 \pm 5.2$ vs. $+6.6 \pm 4.7\%$ P_{max} , for the NH and HH groups, respectively).

Body fat mass and weight

Body weight (69.5 ± 5.9 vs. 69.6 ± 5.6 kg for the NH group and 69.9 ± 6.4 vs. 69.1 ± 6.2 kg for the HH group) and fat mass percentage (9.9 ± 1.8 vs. $9.1 \pm 1.3\%$ for the NH group and 10.3 ± 1.4 vs. $8.4 \pm 0.7\%$ for the HH group) did not differ between groups.

Night S_pO_2 and heart rate

No differences in average values of night HR were found between the groups or between different days (51 ± 1 and 50 ± 2 bpm, for the NH and HH groups, respectively). Conversely, although mean S_pO_2 values ([Fig. 4B](#)) were similar during the control nights (before the camps, Pre1 and Pre2), they were higher in the NH group than in the HH group between day 1 and day 18 (D1 to D18) (92.1 ± 0.3 vs. 90.9 ± 0.3 , for the NH and HH groups, respectively; $P < 0.001$) and

Table 1. Physiological parameters before (Pre-) and after (Post-) the Live High-Train Low (LHTL) camps for the normobaric hypoxia (NH) and hypobaric hypoxia (HH) groups.

		Pre-	Post-	Delta %
VO _{2max}	NH	65.4 ± 7.8	69.1 ± 5.6 **	6.1 ± 6.8
[mL.kg ⁻¹ .min ⁻¹]	HH	69.2 ± 8.9	73.2 ± 7.1 **	5.2 ± 4.8
HR _{max}	NH	187 ± 7	188 ± 5	0.6 ± 2.6
[b.min ⁻¹]	HH	185 ± 9	185 ± 8	0.1 ± 1.9
P _{max}	NH	353 ± 43	385 ± 38 ***	9.6 ± 5.2
[W]	HH	378 ± 24	403 ± 32 ***	6.6 ± 4.7
VE _{max}	NH	178.9 ± 17.8	184.3 ± 14.6	3.7 ± 10.2
[l.min ⁻¹]	HH	182.6 ± 34	188.1 ± 19.3	4.5 ± 9.3

VO_{2max} maximal oxygen uptake; HR_{max} maximal heart rate; P_{max} maximal power output; VE_{max} maximal ventilation. Data are mean ± SD; **P<0.01 and ***P<0.001 for differences between Pre- and Post-.

doi:10.1371/journal.pone.0114418.t001

remained higher (P<0.05) during each of the two nights following the camps (Post1: 94.7 ± 0.5 vs. 93.5 ± 0.9% and Post2: 94.8 ± 0.6 vs. 93.7 ± 1.3%).

Breathing frequency

The average values of breathing frequency were similar between the groups during the two nights prior to the camp (14.0 ± 1.8 and 13.9 ± 1.5 breath.min⁻¹ for NH and HH, respectively). However, breathing frequencies were lower in the NH group than in the HH group (13.9 ± 2.1 vs. 15.5 ± 1.5 breath.min⁻¹, P<0.05) during the LHTL camp (D1 to D18) and remained lower upon the camp's completion (13.7 ± 1.9 vs. 15.1 ± 1.3 breath.min⁻¹, P<0.05).

Total Haemoglobin mass

All results are presented in [Table 2](#). Both groups increased their total haemoglobin masses during the study period (912 ± 96 vs. 936 ± 103 g and 950 ± 115 vs. 967 ± 122 g for the NH and HH groups, respectively, P<0.001).

Blood Parameters

All blood parameters are presented in [Table 2](#). The RBC number, [Hb] and Hct were each lower in the NH group than in the HH group following camp. Additionally, the initial ferritin values were not different between groups and were within reference ranges (98.7 ± 75.9 vs. 105.3 ± 51.9 ng/mL for NH and HH, respectively). A larger decrease in [EPO] was noted in the HH group compared with the NH group with return to 1150 m ([table 2](#)).

Hypoxic doses and P_iO₂

The daily (12.2 ± 0.3 vs. 16.8 ± 3.1 h, P<0.001) and total (220.1 ± 0.9 vs. 302.9 ± 5.5 h, P<0.001) hypoxic doses were lower in the NH group than in the




Fig. 4. A. Daily values of inspired pressure of oxygen (P_iO₂ in mmHg) during the Live High-Train Low (LHTL) camps for the normobaric hypoxia (NH) and hypobaric hypoxia (HH) groups. B. Mean values of night oxygen pulse saturation (S_pO₂). Data are presented in mean ± standard error. Pre1-Pre2: measurements before the camps (1150 m, Prémanson, France); D01-D18: measurement during the camps (NH: hypoxic room in Prémanson, France; HH: Fiescheralp, Switzerland). *P<0.05, ####P<0.001 for differences between groups.

doi:10.1371/journal.pone.0114418.g004

Table 2. Haematological parameters before (Pre-) and after (Post-) the Live High-Train Low (LHTL) camps for the normobaric hypoxia (NH) and hypobaric hypoxia (HH) groups.

		Pre-	Post-	Delta %
EPO	NH	3.85 ± 1.42	3.37 ± 1.59	-14.1 ± 18.3
[mU/mL]	HH	4.96 ± 3.73	3.92 ± 3.94*	-34.5 ± 27.5 [#]
RBC	NH	5.26 ± 0.39	5.08 ± 0.46	-3.5 ± 4.8
[u/μl]	HH	5.15 ± 0.36	5.18 ± 0.43 ^{##}	0.6 ± 3.8 [#]
HGB	NH	15.75 ± 1.07	15.75 ± 1.14	0.1 ± 5.1
[g/dl]	HH	15.53 ± 0.88	16.18 ± 1.08 [#]	4.2 ± 3.9 [#]
Hct	NH	46.25 ± 2.89	45.19 ± 2.97	-2.2 ± 5.6
[%]	HH	45.24 ± 2.43	46.33 ± 2.62 [#]	2.5 ± 3.8 [#]
MCV	NH	88.06 ± 3.71	89.22 ± 3.16 **	1.4 ± 1.3
[fl]	HH	88.06 ± 4.74	89.69 ± 4.51 ***	1.9 ± 1.4
MCH	NH	29.96 ± 1.13	31.09 ± 1.27 ***	3.8 ± 0.9
[pg]	HH	30.22 ± 1.27	31.32 ± 1.34 ***	3.6 ± 1.2
MCHC	NH	34.03 ± 0.72	34.85 ± 0.58 ***	2.4 ± 1.2
[g/dl]	HH	34.33 ± 1.01	34.92 ± 0.79 ***	1.7 ± 2.0
RET	NH	0.89 ± 0.31	1.03 ± 0.28 **	21.6 ± 30.0
[%]	HH	0.98 ± 0.24	1.18 ± 0.38 [#]	23.2 ± 34.1 [#]
IRF	NH	6.72 ± 3.61	5.37 ± 2.17	-9.02 ± 33.95
[%]	HH	6.06 ± 1.77	4.76 ± 2.75	-19.09 ± 36.46
Hb _{mass}	NH	912.4 ± 96.6	935.9 ± 102.6 ***	2.6 ± 1.9
[g]	HH	946.8 ± 126.7	978.6 ± 131.6 ***	3.4 ± 2.1
RCV	NH	2675.8 ± 295.4	2692.1 ± 289.9	0.7 ± 2.8
[ml]	HH	2734.1 ± 306.5	2778.7 ± 324.1	1.64 ± 3.1
BV	NH	6358.3 ± 583.8	6553.6 ± 664.1	3.1 ± 4.8
[ml]	HH	6617.1 ± 744.1	6557.5 ± 821.4	-1.0 ± 4.2 ^{##}
PV	NH	3682.6 ± 384.6	3861.6 ± 460.9	5.1 ± 8.6
[ml]	HH	3883.1 ± 505.3	3778.8 ± 551.8	-2.7 ± 6.1 ^{##}

EPO erythropoietin; RBC red blood cells; HGB haemoglobin; Hct hematocrit; MCV mean cell volume; MCH mean cell haemoglobin; MCHC mean cell haemoglobin concentration; RET reticulocytes; IRF immature reticulocyte fraction; Hb_{mass} haemoglobin mass; RCV red cell volume; BV blood volume; PV plasma volume. Data are mean ± SD; *P<0.05, **P<0.01 and ***P<0.001 for differences between Pre- and Post-; [#]P<0.05 and ^{##}P<0.01 for differences between NH and HH.

doi:10.1371/journal.pone.0114418.t002

HH group. No differences were found in either daily or average P_iO₂ values between the two training camps (121.7 ± 1.2 vs. 121.4 ± 0.9 mmHg for the NH and HH groups, respectively, [Fig. 4A](#)).

Questionnaires

The mean Lake Louise Score was 1.2 ± 0.4 for the NH group and 1.1 ± 0.4 for the HH group. No differences were found between the groups. DALDA Part B results were not different between the groups, and scores did not change across days (2.3 ± 0.7 vs. 2.3 ± 0.6 for the NH and HH groups, respectively). DALDA Part A results included a significantly higher score for the NH group than for the HH

group from D07 to D11 and D15 to D16 ($P < 0.05$). The average VAS value for the sleep quality of the entire camp was lower in the NH group (6.0 ± 0.4 vs. 6.4 ± 0.4 for the NH and HH groups, respectively; $P < 0.001$). The GSQSS was significantly higher for the NH group (4.7 ± 1.1 vs. 3.6 ± 0.8 for the NH and HH groups, respectively; $P < 0.001$), indicating poorer sleep quality for the NH group than for the HH group. However, waking state VAS scores were not different between the groups (5.9 ± 0.5 vs. 5.7 ± 0.5 for the NH and HH groups, respectively).

Correlations

A positive correlation was found between the mean Hb_{mass} (in g) and $VO_{2\text{max}}$ ($\text{mL}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$) values of both groups following camp ($r = 0.68$, $P < 0.01$ and $r = 0.86$, $P < 0.001$ for the NH and HH groups, respectively). We did not find any correlations between changes in Hb_{mass} and $VO_{2\text{max}}$ or between initial value of Hb_{mass} and any other parameter.

Discussion

The present study demonstrated that an 18-d LHTL altitude camp performed under either NH or HH conditions induced different physiological and performance responses: (i) during LHTL, longer hypoxic exposure, larger night desaturation levels, and higher breathing frequencies were noted in the HH group; (ii) immediately after LHTL, larger haematological changes occurred in the HH group, but similar increases were observed in Hb_{mass} and $VO_{2\text{max}}$; (iii) finally, larger performance enhancements were measured in the HH group 3 weeks after return to sea level.

Differences in daily responses were found between our experimental groups. The total hypoxic dose was higher for the HH group than for the NH group (300 vs. 220 h). The present study aimed to compare two LHTL altitude training camps (simulated versus real altitude) in “real conditions”, corresponding to those encountered by elite athletes in their training practices. In this context, the daily exposures reported in the present study (12 vs. 17 $\text{h}\cdot\text{d}^{-1}$ in the NH and HH groups, respectively) are consistent with those reported previously with hypoxic exposures of 8 – 12 $\text{h}\cdot\text{d}^{-1}$ [4, 5] and 18 $\text{h}\cdot\text{d}^{-1}$ [7, 19] in NH and HH conditions, respectively. Meta-analysis results suggest that optimal durations are of 11 and 18 $\text{h}\cdot\text{d}^{-1}$ [27]. There is a clear dose-response effect between the hypoxic dose and the haematological responses, as highlighted by Levine and Stray-Gundersen’s study [23]. However, differences in hypoxic doses between the NH and HH groups cannot be easily reduced, as long duration confinement for individuals of the NH group may cause other complications, including detrimental reductions in plasma volume [34]. Aside from the difference in total hypoxic exposure, the HH group also experienced fewer transitions between “high” (2250 m) and “low” altitudes. LHTL under NH conditions implies numerous daily exposures to normoxia and many transitions to and from hypoxic conditions. Therefore, the

exposure is more intermittent than it would be for LHTL under HH conditions (e.g., the number of daily shifts between altitudes was 7 vs. 2 in the NH and HH groups, respectively). Taken together, we assume that these intermittent characteristics cannot be ruled out as explanations for noted altitude training adaptations. In addition, Navarette-Opazo & Mitchell [35] demonstrated in a recent meta-analysis, that the number of cycles per day was one of the most important variables for the efficiency of the intermittent hypoxic methods. Interestingly, in a similar way, Garvican et al. [22] described the “*oscillating nature of LHTL*,” the daily descents to sea level and the associated normoxic exposures as an explanation for the less dramatic fall in [EPO] from Pre- to Post-, compared to changes associated with continuous exposure to high altitudes. This observation may also explain the differences (almost twice as much in the HH group) in [EPO] decreases noted between the two groups.

Night measurements also indicated that the two conditions were not similar. Night arterial oxygen saturations were higher in the NH group than in the HH group, while breathing frequencies were lower (Fig. 4B). Similarly, Savourey et al. [17] first demonstrated that HH induced greater respiratory frequencies, lower tidal volumes, and minute ventilation values over short time periods; thus, suggesting higher amounts of alveolar physiologic dead space, which is associated with ventilatory alkalosis and hypocapnia [14]. Later, similar conclusions have also been reported by Richard and Koehle [13] and Faiss et al. [18]. Changes in fluid balance have also been shown with differences between the HH and NH conditions [15]. Additionally, barometric pressure (PB) modifies fluid circulation and trans-alveoli-capillary membrane flux [36]. This may induce a stronger pulmonary vasoconstriction in the HH group and modify oxygen diffusion by decreasing the pressure gradient [14]. PB may also influence N_2 and O_2 concentrations in cerebrospinal fluid, as well as central regulation of ventilation [12]. In addition to the reported alteration in ventilatory pattern, all these alterations might contribute also to the lower mean sleeping S_pO_2 values in the HH group.

It is interesting to note that these lower S_pO_2 values were maintained after subjects returned to 1150 m during the two nights immediately following the camp. Several studies [17, 37] have reported a more rapid blood desaturation under HH conditions, leading to a longer duration of hypoxemia. This is in line with the present results as we observed a larger decrease in S_pO_2 from the first night under HH conditions and the maintenance of these lower values during the whole LHTL camp. Interestingly, the higher breathing frequencies while sleeping in HH conditions at 2250 m were also maintained during the two nights spent at 1150 m following the camp. This larger desaturation, which was most likely influenced by a lower tidal volume, potentially induced more severe hypoxemia in the HH group and delayed performance enhancement.

Specific short-term post-hypoxic responses were observed immediately following the camps. These responses were noted primarily among haematological parameters and revealed differences between the groups that were clearly influenced by differences in both daily and total hypoxic exposures. To date, there

is consensus on neither the dose-response relationship between hypoxic stimulus and Hb_{mass} increase nor on the recommendation in terms of total hypoxic exposure duration. For instance, Richalet and Gore have recommended an exposure of 216 h [20], Garvican et al. of 300 h [22], and Wilber et al. a minimum of 4 weeks with at least 22 h.d⁻¹ [25]. However, although the increase in Hb_{mass} immediately following camp was significant, the magnitude of the increase was the same in both groups (2.6 vs. 3.4% for the NH and HH groups, respectively). Our results are consistent with those of previous studies in which Hb_{mass} was increased by 3–4% following several LHTL protocols [7, 19, 22, 28, 38] and illustrate an enhanced oxygen transport capacity as a result of an erythropoietic response. These findings are also consistent with the dose-response relationship and correspond to an average rise of 1% per 100 h of exposure [1, 38]. Gore et al. [38] also suggested that the amount of the hypoxic dose or the level of altitude should be higher in cases involving the use of simulated altitudes to produce equivalent results at real altitudes (*i.e.* 3000 m in NH with LHTL corresponding to 2320 m in HH with classical altitude training). Further, our results showed a 5–6% increase in VO_{2max} in both groups following LHTL, an increase that is commonly observed under HH conditions [19] as well as NH conditions [26, 39], although the phenomenon is more common in the former [3, 40]. In the present study, the difference of 80 h (220 vs. 300 h for the NH and HH groups, respectively) is likely the primary cause of the larger increases in Hct, [Hb] and RET in the HH group. These larger increases with no difference in Hb_{mass} increase could also be explained by plasma shift differences.

We also reported differences in plasma and blood volume changes during the study period between the NH and HH groups. Of interest, is that diuresis and changes in fluid balance have been shown to be different between HH and NH (*i.e.* larger diuresis for NH and larger fluid retention for HH) [15, 41]. The influence of PV changes (*e.g.*, expansion and reduction) on performance enhancement are well-known [42]. It is known that plasma volume may increase until at least 16 days following an altitude training camp [43]. One may speculate that the non-significant difference in plasma volume observed at Post- between the two groups would not occur anymore at Post-21, suggesting a potential larger increase in PV for the HH group during these 3 weeks. This potential hemodynamic enhancement would partially explain the longer delay in performance enhancement compared to the NH group. However, based on the existing contradictory literature, it is difficult to speculate on the maintenance of the Hb_{mass} gains at Post-21. Garvican-Lewis et al. [44] reported a 4% increase two weeks after 11 days of LHTL under NH conditions at 3000 m (14 h.d-1). However, a persistent increase in Hb_{mass} post-exposure does not mean that it did not decrease from the initial elevation during the days following the exposure. Several studies show a relatively linear decrease in Hb and Hct starting as soon as hypoxic stimulus is removed. For example, Gough et al. [45], reported a drop from post to post-14; and Garvican et al. [22], demonstrated that Hb_{mass} had started to drop off by Post-4 and was no different from control after 10 days at sea

level. Similarly, elite altitude-native Kenyan runners showed a significant 20 g decrease after 21 days at sea level (Fig. 2A in [46]).

One of the most important findings of the present study is the performance enhancement noted in the HH group three weeks after camp. Our results are consistent with those of the meta-analysis by Bonetti and Hopkins [27], which described a “terrestrial” LHTL protocol that induced additional benefits relative to the performances of elite athletes as estimated by power output increases of 4.0% under HH conditions and 0.6% under NH conditions.

Delayed (in our case, three weeks after LHTL) performance enhancement has been observed in several [7, 26] but not all studies [39]. The following mechanisms have been proposed: enhanced stroke volume compensating for the reduction in heart rate [47], enhanced efficiency [41], and increased VO_2 and power output at the lactic threshold [26]. Recently, Chapman et al. [48] described the following three components, which may influence performance changes following altitude training: the timing of the decay of red cell mass, the consequences of ventilatory acclimatisation, and the alterations in the biomechanical and neuromuscular factors associated with force production. Regarding the first component, we cannot determine if the unknown decay of Hb_{mass} (see above) in the present study influenced the difference in performance enhancement between groups at Post-21. The observed differences in the ventilatory pattern, as evidenced by the higher night breathing frequency in the HH group, could have influenced the delayed performance difference between groups. Finally, it is unlikely that there was any biomechanical alteration in either group who trained “low” at 1100–1200 m, as suggested by the preliminary results of Laymon et al. [49]. Therefore, a hypoxic-induced alteration in running style was probably not involved in the observed difference at Post-21.

Strengths and limitations

Our primary aim was to compare LHTL training camps under real and simulated altitude in an ecological setting (*e.g.*, by reproducing “real life” conditions of daily exposures and camp durations as described in previous LHTL studies under NH and HH conditions, respectively) rather than investigating the efficiency of LHTL, which has previously been documented. For this reason, we did not include a sea level control group. To the best of our knowledge, our study is the first to report differences in performance enhancement following direct comparisons of prolonged altitude training under NH and HH conditions. The athletes were well trained as shown by their training loads, $\text{VO}_{2\text{max}}$, and performance levels. The groups were matched according to the $\text{VO}_{2\text{max}}$ values. Additionally, the athletes’ training load and content were quantified and matched during the 6-month period before the study, which included a suitable lead-in period. To our knowledge, this study is the first where training loads and altitude levels were entirely matched on a daily basis during the entire LHTL period. The current study emphasizes the importance of well-controlled studies to achieve a better understanding of the mechanisms and potential benefits of altitude training [50].

The primary limitation of this study was that no measurements of total haemoglobin mass or other haematological parameters were completed at one or three weeks following LHTL due to logistical constraints. In addition, since our aim was to compare two typical 18-d LHTL camps in “real” conditions, the hypoxic doses were different. One cannot rule out that the physiological and performance responses would be less dissimilar between NH and HH with a close matching of hypoxic doses.

Perspectives

This study questions the relationship between modes of prolonged hypoxic exposure and subsequent performance improvement. Real altitude conditions (HH) were more demanding than the simulated altitude (NH) utilised in training camps of the same duration. However, in general, hypoxic chambers make adjustments possible and continue to attract interest because of their practicality. Chapman [51] emphasises that the response to training and competition at high altitudes is individual, and that timing the return to competition after altitude training must also be individualised to obtain optimal sea level performance [48]. Normobaric hypoxia devices offer these individualisation possibilities in terms of hypoxic doses and altitude adjustments. Finally, further studies are necessary to assess the physiological responses of these hypoxic training methods to equivalent hypoxic doses.

This study highlights the different physiological adaptations noted in the HH and NH LHTL camps. Our results suggest that future investigations should increase the altitude of the normobaric hypoxia group to reach the same level of desaturation as that experienced under hypobaric hypoxic conditions and lengthen the durations of the camps to obtain hypoxic doses similar to those experienced under hypobaric hypoxic conditions.

Conclusion

The primary finding of the study is that there were significant differences in the responses to a LHTL training camp in NH compared to HH. Specifically, our results included greater performance enhancements in the HH group three weeks after LHTL, greater significance in haematological changes within the HH group following camp, greater night desaturation levels, and higher breathing frequencies in the HH group, with similar increases in Hb_{mass} and VO_{2max} following LHTL in both NH and HH. Additionally, one cannot rule out other factors, including differences in sleep quality, desaturation level, breathing patterns, fewer transitions between high and low altitudes (*e.g.*, intermittence) or different responses relative to plasma volumes and [EPO] following camp.

Acknowledgments

This study was funded by grants from the Ministère des Sports, de la Jeunesse, de l'Éducation Populaire et de la Vie Associative (MSJEPVA)/Institut National du Sport, de l'Expertise et de la Performance (INSEP) and from Bundesamt für Sport (BASPO; Switzerland). The authors thank the director Vincent Jacquet and the staff of the National Ski-Nordic Centre (Prémanon, France) as well as the director Claudio Rossetti and the staff of the Fierendorf Center (Fiesch, Switzerland) for their invaluable assistance and access to facilities. The authors also thank Mattia Bertsch for the use of the instrumented t-shirt and Olivier Girard for the final revision of the manuscript. The authors have no conflicts of interest, source of funding or financial ties to disclose and no current or past relationship with companies or manufacturers who could benefit from the results of the present study.

Author Contributions

Conceived and designed the experiments: JJS LS RF AH JPW GPM. Performed the experiments: JJS LS RC RF AH JPW BR AD NR GPM. Analyzed the data: JJS LS AH JPW NR GPM. Contributed reagents/materials/analysis tools: LS JPW NR GPM. Wrote the paper: JJS GPM.

References

1. Clark SA, Quod MJ, Clark MA, Martin DT, Saunders PU, et al. (2009) Time course of haemoglobin mass during 21 days live high:train low simulated altitude. *Eur J Appl Physiol* 106: 399–406.
2. Gore CJ, Hahn AG, Aughey RJ, Martin DT, Ashenden MJ, et al. (2001) Live high:train low increases muscle buffer capacity and submaximal cycling efficiency. *Acta Physiol Scand* 173: 275–286.
3. Hahn AG, Gore CJ, Martin DT, Ashenden MJ, Roberts AD, et al. (2001) An evaluation of the concept of living at moderate altitude and training at sea level. *Comp Biochem Physiol A Mol Integr Physiol* 128: 777–789.
4. Roberts AD, Clark SA, Townsend NE, Anderson ME, Gore CJ, et al. (2003) Changes in performance, maximal oxygen uptake and maximal accumulated oxygen deficit after 5, 10 and 15 days of live high:train low altitude exposure. *Eur J Appl Physiol* 88: 390–395.
5. Saunders PU, Telford RD, Pyne DB, Cunningham RB, Gore CJ, et al. (2004) Improved running economy in elite runners after 20 days of simulated moderate-altitude exposure. *J Appl Physiol* (1985) 96: 931–937.
6. Dehnert C, Hutler M, Liu Y, Menold E, Netzer C, et al. (2002) Erythropoiesis and performance after two weeks of living high and training low in well trained triathletes. *Int J Sports Med* 23: 561–566.
7. Levine BD, Stray-Gundersen J (1997) "Living high-training low": effect of moderate-altitude acclimatization with low-altitude training on performance. *J Appl Physiol* (1985) 83: 102–112.
8. Stray-Gundersen J, Levine BD (2008) Live high, train low at natural altitude. *Scand J Med Sci Sports* 18 Suppl 1: 21–28.
9. Wehrli JP, Zuest P, Hallen J, Marti B (2006) Live high-train low for 24 days increases hemoglobin mass and red cell volume in elite endurance athletes. *J Appl Physiol* (1985) 100: 1938–1945.
10. Millet GP, Roels B, Schmitt L, Woorons X, Richalet JP (2010) Combining hypoxic methods for peak performance. *Sports Med* 40: 1–25.

11. **Mounier R, Brugniaux JV** (2012) Counterpoint: Hypobaric hypoxia does not induce different responses from normobaric hypoxia. *J Appl Physiol* (1985) 112: 1784–1786.
12. **Conkin J, Wessel JH, 3rd** (2008) Critique of the equivalent air altitude model. *Aviat Space Environ Med* 79: 975–982.
13. **Richard NA, Koehle MS** (2012) Differences in cardio-ventilatory responses to hypobaric and normobaric hypoxia: a review. *Aviat Space Environ Med* 83: 677–684.
14. **Millet GP, Faiss R, Pialoux V** (2012) Point: Hypobaric hypoxia induces different physiological responses from normobaric hypoxia. *J Appl Physiol* (1985) 112: 1783–1784.
15. **Loeppky JA, Roach RC, Maes D, Hinghofer-Szalkay H, Roessler A, et al.** (2005) Role of hypobaria in fluid balance response to hypoxia. *High Alt Med Biol* 6: 60–71.
16. **Fulco CS, Muza SR, Beidleman BA, Demes R, Staab JE, et al.** (2011) Effect of repeated normobaric hypoxia exposures during sleep on acute mountain sickness, exercise performance, and sleep during exposure to terrestrial altitude. *Am J Physiol Regul Integr Comp Physiol* 300: R428–436.
17. **Savourey G, Launay JC, Besnard Y, Guinet A, Travers S** (2003) Normo- and hypobaric hypoxia: are there any physiological differences? *Eur J Appl Physiol* 89: 122–126.
18. **Faiss R, Pialoux V, Sartori C, Faes C, Deriaz O, et al.** (2013) Ventilation, oxidative stress, and nitric oxide in hypobaric versus normobaric hypoxia. *Med Sci Sports Exerc* 45: 253–260.
19. **Wehrlin JP, Marti B** (2006) Live high-train low associated with increased haemoglobin mass as preparation for the 2003 World Championships in two native European world class runners. *Br J Sports Med* 40: e3; discussion e3.
20. **Richalet JP, Gore CJ** (2008) Live and/or sleep high:train low, using normobaric hypoxia. *Scand J Med Sci Sports* 18 Suppl 1: 29–37.
21. **Rusko HK, Tikkanen HO, Peltonen JE** (2004) Altitude and endurance training. *J Sports Sci* 22: 928–944; discussion 945.
22. **Garvican L, Martin D, Quod M, Stephens B, Sassi A, et al.** (2012) Time course of the hemoglobin mass response to natural altitude training in elite endurance cyclists. *Scand J Med Sci Sports* 22: 95–103.
23. **Levine BD, Stray-Gundersen J** (2006) Dose-response of altitude training: how much altitude is enough? *Adv Exp Med Biol* 588: 233–247.
24. **Stray-Gundersen J, Chapman RF, Levine BD** (2001) "Living high-training low" altitude training improves sea level performance in male and female elite runners. *J Appl Physiol* (1985) 91: 1113–1120.
25. **Wilber RL, Stray-Gundersen J, Levine BD** (2007) Effect of hypoxic "dose" on physiological responses and sea-level performance. *Med Sci Sports Exerc* 39: 1590–1599.
26. **Schmitt L, Millet G, Robach P, Nicolet G, Brugniaux JV, et al.** (2006) Influence of "living high-training low" on aerobic performance and economy of work in elite athletes. *Eur J Appl Physiol* 97: 627–636.
27. **Bonetti DL, Hopkins WG** (2009) Sea-Level Exercise Performance Following Adaptation to Hypoxia: A Meta-Analysis. *Sports Med* 39: 107–127.
28. **Robertson EY, Saunders PU, Pyne DB, Gore CJ, Anson JM** (2010) Effectiveness of intermittent training in hypoxia combined with live high/train low. *Eur J Appl Physiol* 110: 379–387.
29. **Chapman RF, Karlisen T, Resaland GK, Ge RL, Harber MP, et al.** (2014) Defining the "dose" of altitude training: how high to live for optimal sea level performance enhancement. *J Appl Physiol* (1985) 116: 595–603.
30. **Cejuela Anta R, Esteve-Lanao J** (2011) Training load quantification in triathlon. *Journal of human Sport & Exercise* 6: 218–232.
31. **Steiner T, Wehrlin JP** (2011) Does hemoglobin mass increase from age 16 to 21 and 28 in elite endurance athletes? *Med Sci Sports Exerc* 43: 1735–1743.
32. **Schmidt W, Prommer N** (2005) The optimised CO-rebreathing method: a new tool to determine total haemoglobin mass routinely. *Eur J Appl Physiol* 95: 486–495.
33. **Lamon S, Robinson N, Saugy M** (2010) Procedures for monitoring recombinant erythropoietin and analogs in doping. *Endocrinol Metab Clin North Am* 39: 141–154, x.

34. **Siebenmann C, Robach P, Jacobs RA, Rasmussen P, Nordsborg N, et al.** (2012) "Live high-train low" using normobaric hypoxia: a double-blinded, placebo-controlled study. *J Appl Physiol* (1985) 112: 106–117.
35. **Navarrete-Opazo AA, Mitchell GS** (2014) Therapeutic Potential of Intermittent Hypoxia: A Matter of Dose. *Am J Physiol Regul Integr Comp Physiol*.
36. **Levine BD, Kubo K, Kobayashi T, Fukushima M, Shibamoto T, et al.** (1988) Role of barometric pressure in pulmonary fluid balance and oxygen transport. *J Appl Physiol* (1985) 64: 419–428.
37. **Singh B, Cable GG, Hampson GV, Pascoe GD, Corbett M, et al.** (2010) Hypoxia awareness training for aircrew: a comparison of two techniques. *Aviat Space Environ Med* 81: 857–863.
38. **Gore CJ, Sharpe K, Garvican-Lewis LA, Saunders PU, Humberstone CE, et al.** (2013) Altitude training and haemoglobin mass from the optimised carbon monoxide rebreathing method determined by a meta-analysis. *Br J Sports Med* 47 Suppl 1: i31–i39.
39. **Robach P, Schmitt L, Brugniaux JV, Roels B, Millet G, et al.** (2006) Living high-training low: effect on erythropoiesis and aerobic performance in highly-trained swimmers. *Eur J Appl Physiol* 96: 423–433.
40. **Ashenden MJ, Gore CJ, Parisotto R, Sharpe K, Hopkins WG, et al.** (2003) Effect of altitude on second-generation blood tests to detect erythropoietin abuse by athletes. *Haematologica* 88: 1053–1062.
41. **Gore CJ, Clark SA, Saunders PU** (2007) Nonhematological mechanisms of improved sea-level performance after hypoxic exposure. *Med Sci Sports Exerc* 39: 1600–1609.
42. **Coyle EF, Hopper MK, Coggan AR** (1990) Maximal oxygen uptake relative to plasma volume expansion. *Int J Sports Med* 11: 116–119.
43. **Heinicke K, Wolfarth B, Winchenbach P, Biermann B, Schmid A, et al.** (2001) Blood volume and hemoglobin mass in elite athletes of different disciplines. *International journal of sports medicine* 22: 504–512.
44. **Garvican-Lewis LA, Clark SA, Polglaze T, McFadden G, Gore CJ** (2013) Ten days of simulated live high:train low altitude training increases Hbmass in elite water polo players. *Br J Sports Med* 47 Suppl 1: i70–73.
45. **Gough CE, Saunders PU, Fowlie J, Savage B, Pyne DB, et al.** (2012) Influence of altitude training modality on performance and total haemoglobin mass in elite swimmers. *Eur J Appl Physiol* 112: 3275–3285.
46. **Prommer N, Thoma S, Quecke L, Gutekunst T, Volzke C, et al.** (2010) Total hemoglobin mass and blood volume of elite Kenyan runners. *Med Sci Sports Exerc* 42: 791–797.
47. **Burtscher M, Nachbauer W, Baumgartl P, Philadelphia M** (1996) Benefits of training at moderate altitude versus sea level training in amateur runners. *Eur J Appl Physiol Occup Physiol* 74: 558–563.
48. **Chapman RF, Laymon Stickford AS, Lundby C, Levine BD** (2013) Timing of Return From Altitude Training for Optimal Sea Level Performance. *J Appl Physiol* (1985) 116: 837–843.
49. **Laymon ASL EA, Mckenzie JM, Wilhite DP, Chapman RF** (2009) Running economy changes after altitude training: role of running mechanics. *Medicine and science in sports and exercise* 41: 320.
50. **Lundby C, Millet GP, Calbet JA, Bartsch P, Subudhi AW** (2012) Does 'altitude training' increase exercise performance in elite athletes? *Br J Sports Med* 46: 792–795.
51. **Chapman RF** (2013) The individual response to training and competition at altitude. *Br J Sports Med* 47 Suppl 1: i40–44.