

Evaluation of estimation methods for fitting the three-parameter Weibull distribution to European beech forests

Živa Bončina⁽¹⁻²⁾,
Vasilije Trifković⁽¹⁾,
Christian Rosset⁽³⁾,
Matija Klopčič⁽¹⁾

We evaluated three estimation methods for fitting the three-parameter Weibull distribution to even-aged European beech (*Fagus sylvatica*) forests by using experimental tree diameter data collected in 3709 sample plots (500 m² each). The maximum likelihood estimation method (MLE), the method of moments (MOM) and the method of modified moments type 1 (MM1) were applied for fitting the Weibull function. The goodness-of-fit of stand parameters (total tree number, stand basal area, dominant stand diameter and mean quadratic diameter) was tested by MAE and RMSE, and the probability and cumulative density functions of trees per 5 cm diameter classes were additionally tested by the Kolmogorov-Smirnov test and compared with Kolmogorov-Smirnov's D statistic. All three methods are suitable for estimating stand parameters based on the fitted Weibull function. Fitting the diameter distribution per 5 cm diameter classes at the plot level was less accurate due to the low number of trees or irregular diameter distribution of trees. The MM1 method was found to be the most suitable for fitting the three-parameter Weibull distribution to beech forests represented by data derived from small plots.

Keywords: Diameter Distribution Fitting, Weibull Function, Parameter Estimation, Inventory Data, Circular Sample Plots, Near-natural Forests, *Fagus Sylvatica*, Slovenia

Introduction

Tree diameter distribution is a key piece of information about forest stand structure (Pretzsch 2009) which forest managers use to evaluate recent forest management and make decisions about further silvicultural treatments (Burkhart & Tomé 2012, Bassil et al. 2019). Tree diameter distribution is integrated into several individual-tree and stand growth models (Schütz & Rosset 2020, Pretzsch et al. 2002, Hasenauer et al. 2006, Sterba et al. 1995). In stand models, parameters of tree diameter distribution are often determined by regressions (Burkhart & Tomé 2012) in which easily measured stand parameters, such as stand basal area (BA), mean stand diameter and total number of trees (N), are used as explanatory variables (Nord-Larsen & Cao

2006, Schütz & Rosset 2020).

The tree diameter distribution of forest stands is described either by the frequency of trees per diameter class or by estimated parameters of the theoretical distribution (Kangas & Maltamo 2006). Theoretical tree diameter distributions can be defined based on the number of trees per diameter class (Cao 2004) or the BA of trees per diameter class (Kangas & Maltamo 2000, Siipilehto 1999). Fitting the diameter distribution with the BA of trees provides better access to the right part of the distribution, representing large-sized (and more valuable) trees, while fitting with N provides better fitting at the left part of distribution (i.e., small-sized trees).

Several distribution functions have been used for fitting diameter distribution to for-

est stands, including exponential, beta, normal, lognormal, Johnson's SB, Weibull and others (Palahi et al. 2007). Since its introduction by Bailey & Dell (1973), the Weibull distribution (Weibull 1939) has been widely used in forest science due to its flexibility, the relative simplicity of parameter estimation, and its closed cumulative density functional form. The Weibull distribution function describes a wide range of unimodal distributions from reversed-J shaped to exponential and normal (Bailey & Dell 1973). Its special feature is the independency of its parameters, at least at the particular position of the 63.2th percentile (Dubey 1967, Zanakis 1979). Since this percentile strongly correlates with the quadratic mean diameter (QMD) of monospecific and regular stands as an easily obtained stand parameter (Schütz & Rosset 2020), the Weibull function has often been applied for fitting diameter distribution to forest stands. However, Weibull distribution has often been applied in plantations or densely planted forest stands (Bailey & Dell 1973, Cao 2004, Lei 2008, Schmidt et al. 2020) and much less frequently in natural forests. In Europe, the Weibull distribution has been fitted for pure (Rosset et al. 2018, Nord-Larsen & Cao 2006, Schütz & Rosset 2020) and mixed stands (Merganič & Sterba 2006, Siipilehto 1999).

Different estimation methods can be applied to fit the Weibull distribution (Zhang et al. 2003, Lei 2008, Hossain & Zimmer 2003). Maximum likelihood estimation

□ (1) Department of Forestry and Renewable Forest Resources, Biotechnical Faculty, University of Ljubljana, Večna pot 83, 1000 Ljubljana (Slovenia); (2) Slovenia Forest Service, Večna pot 2, 1000 Ljubljana (Slovenia); (3) Forest Ecosystem and Management, School of Agricultural, Forest and Food Sciences, HAFL, Bern University of Applied Sciences, Länggasse 85, 3052 Zollikofen (Switzerland)

@ Matija Klopčič (matija.klopcic@bf.uni-lj.si)

Received: May 30, 2022 - Accepted: Sep 27, 2022

Citation: Bončina Z, Trifković V, Rosset C, Klopčič M (2022). Evaluation of estimation methods for fitting the three-parameter Weibull distribution to European beech forests. *iForest* 15: 484-490. - doi: [10.3832/ifor4145-015](https://doi.org/10.3832/ifor4145-015) [online 2022-12-01]

Communicated by: Andrea Cutini




Fig. 1 - Location of sample plots (black dots, n = 3709) in beech forests of Slovenia.

to beech forests; and (iii) to assess the suitability of small circular plot inventory data for fitting diameter distribution to the Weibull function.

Material and methods

We used data from the permanent sampling plots (n ≈ 98,000) of the Slovenia Forest Service (SFS 2021), which are distributed systematically across the forest area of Slovenia. The plots are circular and concentric; trees of 10-29 cm in diameter at breast height (DBH) were measured in 200 m² plots (radius = 7.98 m), and trees of DBH ≥ 30 cm in 500 m² plots (radius = 12.62 m). To scale up the number of trees to 1 ha, trees within the respective circles were weighted with a representation factor of 50 or 20. In order to study even-aged beech stands only, we included plots in which the share of beech exceeded 80% of the total BA. Plots with fewer than 10 trees were discarded (Poudel & Cao 2013, Palahi et al. 2006, Zhang et al. 2003). The Gini index is a measure of the structural diversity of forest stands (Duduman 2011, Klopčič & Bončina 2011) and ranges from 0 to 1, where higher values indicate more heterogeneous forest stand structure. We analyzed plots with even-aged stand structure, defined by Gini index < 0.33. The Gini index was calculated using the basal area of individual trees on each plot (Klopčič & Bončina 2011). To divide even-sized (even-aged) and uneven-sized stands, k-means cluster analysis was performed, resulting in two clusters divided at the given Gini index of 0.33. To validate such classification, the Sokal Micherer measure of similarity was calculated between the two clusters based on Gini index and the field assessment of stand heterogeneity. The McNemara χ² test indicated a significant similarity between clusters (p < 0.05); therefore, we adopted 0.33 as an adequate threshold. This was in accordance with the literature, as Duduman (2011) defined this threshold at 0.35 and O’Hara et al. (2007) at 0.3.

Beech is the main tree species in Slovenian forests, representing 33% of the total growing stock (SFS 2021). The total area of beech forests amounts to approximately 7500 km², representing 70% of the potential natural forest vegetation in Slovenia (Fig. 1). They are classified into several beech forest types (Bončina et al. 2021). The elevation of beech plots ranges from 161 to 1655 m a.s.l. Even-aged beech stands are characterized by relatively high stand BA and large quadratic mean diameter (QMD) and dominant diameter (DDOM – Tab. 1).

The three-parameter Weibull function (Weibull 1939) was used to characterize the diameter distribution. The probability density function (PDF) has the following form (eqn. 1):

$$f(x) = \left(\frac{C}{B}\right) \left(x - \frac{A}{B}\right)^{C-1} \exp\left[-\left(x - \frac{A}{B}\right)^C\right] \quad (1)$$

where A, B and C are the location, scale and

(MLE) is the most common method used in forestry (Schütz & Rosset 2020, Schmidt et al. 2020), while the method of moments (MOM – Lei 2008) or Bayes estimation (Teimouri & Podlaski 2022) have been applied rather less frequently. For estimating parameters of diameter distribution functions, which is the first step in developing a stand model (Schmidt et al. 2020), it is important to employ an adequate estimation method. However, studies on evaluating estimation methods for fitting the Weibull distribution are quite limited (Zhang et al. 2003, Lei 2008, Gorgoso-Varela et al. 2012, Teimouri et al. 2013, Akram & Hayat 2014), and clear procedures and criteria for defining the most appropriate method are lacking.

Estimation methods for fitting Weibull distributions have mainly been evaluated based on generated data (Nanos & Sjöstedt De Luna 2017, Hossain & Zimmer 2003, Chu & Ke 2012, Teimouri et al. 2013, Pobočková & Sedláčková 2014, Akram & Hayat 2014), while comparative studies based on empirical data are less frequent. They have been conducted in pine plantations (Lei 2008, Teimouri et al. 2013), spruce-fir for-

ests (Zhang et al. 2003) and pine forests (Gorgoso-Varela et al. 2012), but comparative studies in broadleaved forests are lacking. European beech (*Fagus sylvatica* L., hereafter beech) is one of the most common trees species in Central Europe (Bohn et al. 2000). To our knowledge there is no evidence regarding which estimation method is the most appropriate for fitting the Weibull distribution function to beech forests. Various data sources can be applied for fitting the Weibull function, national forest inventories (NFI) being only one of them. In most European countries, the NFI is an important source of tree data, enabling its possible application to stand modeling (Kangas & Maltamo 2006). However, little is known about whether data from small sampling plots are appropriate for the parametrization of the Weibull function (Palahi et al. 2007, Siipilehto 1999). Therefore, the main objectives of our study were (i) to fit the three-parameter Weibull distribution function to even-aged beech stands based on data from small sampling plots by using three estimation methods; (ii) to estimate the adequacy of the three estimation methods for fitting the Weibull distribution

Tab. 1 - Stand characteristics of even-aged beech stands (n = 3709). (SD): standard deviation.

Parameter	Label	Mean	SD	Min	Max
Number of trees per hectare	N	540.69	207.81	230	1320
Number of trees per plot	N _{PL}	16.02	4.48	10	45
Minimum diameter (cm)	D _{MIN}	11.92	3.55	10	25
Maximum diameter (cm)	D _{MAX}	51.57	8.92	20	90
Dominant stand diameter (cm)	D _{DOM}	42.05	6.79	15.00	65.50
Mean quadratic diameter (cm)	QMD	29.07	5.32	12.78	47.20
Stand basal area (m ² ha ⁻¹)	BA	33.73	9.43	4.64	70.68
Gini index of heterogeneity	Gini	0.26	0.05	0.00	0.33

shape parameters, respectively, and x is the independent variable, *i.e.*, DBH in our study. The location parameter A identifies the lower bound of the diameter distribution, *i.e.*, the minimum diameter (Bailey & Dell 1973). When the distribution starts at zero, the three-parameter PDF function becomes a two-parameter PDF function. The scale parameter B corresponds to the 63.2th percentile of the distribution (Schütz & Rosset 2020), while parameter C defines the shape of the Weibull distribution (Bailey & Dell 1973). When $C < 1$ the distribution is reversed-J-shaped, and when $1 < C < 3.6$ the distribution is mound-shaped and positively skewed. When $C \approx 3.6$ it is approximately equivalent to the normal distribution, and when $C > 3.6$ the distribution becomes progressively negatively skewed. The average skewness of all plots ($n = 3709$) was 0.5, ranging from -1.0 to 4.3, while the average kurtosis was -0.3, ranging from -1.9 to 19.9.

Parameters A , B and C were estimated using three estimation methods: the method of maximum likelihood (MLE), the method of moments (MOM) and the method of modified moments type 1 (MM1) using the R package “ForestFit” (Teimouri 2020). For each plot, the parameters were iteratively searched. The estimation methods do not always converge (Akram & Hayat 2014). In order to detect non-convergence of estimates, we cleaned extremely deviated values of parameters A , B and C and predicted BA and N with the method for detecting outliers using the interquartile range (Senthamarai Kannan et al. 2015 – eqn. 2):

$$q_1(x) - 1.5IQR(x) < x < q_3(x) + 1.5IQR(x) \quad (2)$$

where x is the parameter under observation, q_1 and q_3 are the first and third quartiles of the distribution of x and IQR is the interquartile range of parameter x . In total, almost 42% of the plots were excluded from the analysis due to the outlier rule. Of this proportion, 28% were excluded due to the extreme deviation of parameters A , B and C and 14% due to extremely deviated BA or N predictions. Most of these plots were excluded due to deviated values obtained by the MLE method. The percentages of excluded plots due to deviated values of Weibull function parameters were 24%, 6% and 7% of the total plot number for the MLE, MOM and MM1 methods, respectively. The final dataset encompassed 3709 sampling plots (Fig. 1).

The three estimation methods for fitting the Weibull function were evaluated based on: (i) the number of excluded plots due to deviated values; (ii) comparison of the observed and predicted values of stand parameters; and (iii) comparison of the observed and predicted values of parameters within DBH classes.

Stand parameters (*i.e.*, N, BA, DDOM and QMD) for the observed and fitted distributions were compared using the mean absolute error (MAE) and the root-mean-square

Tab. 2 - Descriptive statistics of the Weibull function parameters estimated by the three methods. (SD): standard deviation.

Parameter	Method	Mean	SD	Min	Max
A	MLE	14.04	3.62	8.09	27.50
	MOM	20.99	4.02	11.62	32.40
	MM1	14.25	3.56	10.78	27.50
B	MLE	12.57	4.31	0.14	27.62
	MOM	6.36	1.96	0.28	13.77
	MM1	13.85	4.88	0.04	27.56
C	MLE	0.86	0.47	0.19	2.07
	MOM	1.08	0.08	0.61	1.33
	MM1	1.28	0.24	0.33	2.06

error (RMSE). Since parameter C defines the shape of distribution, we analysed the Pearson’s correlation between the shape parameter C and the Gini index as a measure of stand homogeneity. To test for differences between the fitted and observed diameter distributions for the 5 cm DBH classes, several tests were used: the Kolmogorov-Smirnov (KS) goodness-of-fit test by using the cumulative density function (CDF); KS’s D statistic, which is the maximum difference between empirical and fitted CDF (Zuur et al. 2007); and the MAE and RMSE, both based on the PDF for each 5 cm DBH class by using relative frequencies of trees per DBH class, which enabled comparison of plots regardless of the number of trees. Finally, we calculated the mean values of MAE, RMSE and KS’s D statistic for each estimation method. Estimated Weibull distributions from the three analysed methods were illustrated with three plots showing typical distribution: approximately normal, right skewed (reversed-J-shaped) and multimodal distribution. Since stand structure may affect fitting (Merganič & Sterba 2006), we explored the correlation between stand parameters (BA and QMD) and the mean MAE within DBH classes.

Results

The mean values of the Weibull function

parameters A and B were quite similar when obtained by the MM1 or MLE method, while MOM gave higher values of parameter A and lower values of parameter B (Tab. 2). The mean values of parameter C , indicating the shape of distribution, exhibited the expected values above 1 when obtained by the MM1 and MOM methods, but not by the MLE method. Pearson’s correlation coefficients between the shape parameter C and the Gini index were -0.130, -0.205 and -0.214 for the MLE, MOM and MM1 methods, respectively (all $p < 0.001$).

The differences in N, BA, DDOM and QMD between observed and fitted distributions were quite low for most plots (Tab. 3). The MAE varied between 0.04 and 10.8 m² ha⁻¹ and 3 to 42 trees ha⁻¹ for BA and N, respectively, when different estimation methods were considered. Differences between observed and fitted values were lower for QMD than for DDOM for all methods. Of all methods, MM1 had the smallest differences between the observed and predicted stand parameters.

The MM1 method also outperformed the other two methods in goodness-of-fit measures for distributions per DBH class (Tab. 4). When MAE or RMSE were considered, the performance of the MLE method was similar to that of the MM1 method, while the MOM method performed much poorer (Fig. 2, Tab. 4). However, when the MLE

Tab. 3 - Goodness-of-fit measures for number of trees (N), stand basal area (BA), quadratic mean diameter (QMD) and dominant stand diameter (D_{DOM}) for the three estimation methods ($n = 3709$).

Parameter	Statistics	Method		
		MLE	MOM	MM1
N	MAE (ha ⁻¹)	41.91	21.59	2.51
	RMSE (ha ⁻¹)	82.17	36.48	19.51
BA	MAE (m ² ha ⁻¹)	10.72	2.96	0.04
	RMSE (m ² ha ⁻¹)	15.40	3.31	0.03
QMD	MAE (cm)	6.20	1.28	0.03
	RMSE (cm)	9.05	1.41	0.08
DDOM	MAE (cm)	6.83	7.73	3.25
	RMSE (cm)	9.40	8.14	4.12




Fig. 2 - Ranking of three estimation methods in regard to their performance of fitting the diameter distribution measured by MAE per 5 cm DBH classes ($n = 3709$). Ranks: the method exhibits the lowest MAE (rank 1), the second lowest MAE (rank 2) and the highest MAE (rank 3).

Tab. 4 - Goodness-of-fit measures for diameter distributions per 5 cm DBH classes for the three estimation methods. “Different distributions” is the share (%) of plots with significantly different distributions, detected by the KS test ($p > 0.05$).

Method	Different distributions (%)	Mean MAE	Mean RMSE	Mean KS's D statistic
MLE	3.9	0.086	0.117	0.388
MOM	0.0	0.134	0.195	0.308
MM1	0.0	0.084	0.114	0.257

Tab. 5 - Mean MAE values for tree number per 5 cm DBH classes calculated from the observed and fitted distributions using the MM1 method, shown per quadratic mean diameter (QMD) and stand basal area (BA) classes.

BA class (m ²)	QMD class (cm)			
	10-20	20-30	30-40	> 40
0-20	0.168	0.112	-	-
20-30	0.125	0.088	0.096	-
30-40	0.103	0.073	0.082	0.081
40-50	-	0.060	0.064	0.077

method was used, the observed and fitted distributions significantly differed in almost 4% of the plots (KS test, $p < 0.05$).

To illustrate the estimated Weibull distributions by using the three different methods, three plots with typical distributions are displayed in Fig. 3. The best fit was detected for the near-normal distribution; in this plot the observed and fitted distributions using MLE and MM1 differed by only 4% on average across DBH classes. Most of the criteria indicated poor fitting for the multimodal distribution. A statistically significant difference (KS test, $p < 0.05$) between observed and fitted multimodal distributions was detected for MLE only, although the other two methods also poorly fitted the multimodal distribution. However, all methods applied for the multimodal plots appeared to have a better fit to N in comparison to those with a reversed-J-shaped distribution. The MOM method simulated fewer thinner trees in plots with a multimodal distribution in comparison to the other two methods, which is a consequence of a higher value of parameter A (Tab. 2).

For all methods the impact of stand structure on the goodness of the Weibull function fitting was tested (results shown for MM1 only). The comparative analyses of MAE for diameter distributions fitted by MM1 per BA and QMD classes indicated lower MAE in plots with higher BA and QMD (Tab. 5).

Discussion

Our study is one of the first attempts to fit the Weibull distribution function to empirical data representing semi-natural European beech forests. We used data from small concentric circular sampling plots, which is quite rare when fitting the diameter distribution of forest stands (Lei 2008, Schmidt et al. 2020). Our results showed that the three-parameter Weibull function can be fitted to even-aged beech forests using such data. However, the goodness of fit substantially depends on the diameter distribution of empirical data. In stands




Fig. 3 - Examples of fitted distributions for plots with representative distributions: approximately normal distribution (left), right skewed distribution (middle) and multimodal distribution (right). For each plot, fitted distributions from the three methods (MLE, MOM, MM1) and the histogram of observed empirical data are shown.

with a unimodal distribution fitting was mostly successful, while in stands with a multimodal diameter distribution, fitting was poor or even impossible.

Comparison of estimation methods for fitting the Weibull function

We evaluated three estimation methods for fitting the Weibull function by comparing (i) the number of excluded plots due to deviated values, (ii) the observed and predicted values of stand parameters, and (iii) the observed and predicted values of parameters within DBH classes. The proportion of excluded plots due to extremely deviated values was fairly high. The non-convergence of estimates seems to be an important criterion for selecting the most appropriate fitting method. Our results showed that when using the MLE method the number of excluded plots was almost four times larger than that obtained using MM1 or MOM. Akram & Hayat (2014) found that MLE can outperform other methods for estimating the three-parameter Weibull distribution as long as convergence is attained. This result is in accordance with simulation studies (Chu & Ke 2012, Teimouri et al. 2013, Akram & Hayat 2014, Pobočíková & Sedláčková 2014) indicating that other estimation methods outperformed MLE when the number of trees is small and the shape parameter C smaller than one, which was often the case in our study (Tab. 2). The number of excluded plots, meaning the plots in which the MLE method exhibited non-convergence, would likely be much lower if the proportion of plots with unimodal mound-shaped distributions was higher.

The comparative analysis of the observed and fitted values of N , BA , QMD and $DDOM$ indicated that all stand parameters at the plot level are well predicted. Of all methods, the MM1 method exhibited the best fit for all stand parameters, followed by the MOM and MLE methods.

Goodness of fit for diameter distribution per DBH class was much worse in comparison to the goodness of fit for stand parameters (Tab. 3, Tab. 4). The MM1 method was superior to MOM and MLE again considering the mean values of MAE, RMSE and KS's D statistic for diameter distributions per DBH class. The mean MAE for tree number per DBH class, amounting to around 8% of trees in each 5 cm DBH class, was quite comparable for the MM1 and MLE methods. Gorgoso-Varela et al. (2012) reported lower values of goodness-of-fit measures for fitting the Weibull distribution to homogeneous pine stands. However, they used larger sample plots, and only plots with 30 or more trees were included in the analysis.

The MM1 method outperformed the MLE method with respect to the lowest proportion of plots with a significant difference between observed and the fitted diameter distributions. However, the latter should be interpreted with caution since they are

based on the KS test with relatively low frequencies per DBH class.

Suitability of data from small inventory plots for fitting the Weibull distribution function

Our results indicate that forest inventory data obtained on relatively small sampling plots can be used for fitting the Weibull diameter distribution, but with some limitations. The proportion of plots excluded from the analysis due to extremely deviated values of distribution function parameters or modelled BA and N was almost 42%. Akram & Hayat (2014) calculated the non-convergence rate in a different way but still reported a rate of about 44% for sampling plots with 10 trees per plot and shape parameter 1.5. There are several possible reasons for the non-convergence. The small size of the sampling plots is probably the most important. The expectation that the theoretical distribution can be fitted to data from every small plot often containing a low number of trees is simply unrealistic (Nord-Larsen & Cao 2006, Merganič & Sterba 2006), especially if the DBH classes are narrow. In our study only plots with 10 or more trees were included in the analysis. Ten trees has often been recommended as the minimum number of trees for estimating Weibull distribution parameters (Kangas & Maltamo 2000, Palahi et al. 2006, Poudel & Cao 2013), although some studies have suggested 15-30 trees or even more (Merganič & Sterba 2006, Palahi et al. 2007, Gorgoso-Varela et al. 2012, Sghaier et al. 2016, Schütz & Rosset 2020). The size of our sampling plots was 0.05 ha. In similar studies sampling plots of approximately the same size were applied (Zhang et al. 2003), but larger plots were typically used, for instance, > 0.25 ha (Schütz & Rosset 2020) and 0.62 ha (Cao 2004). Due to the threshold value of 10 trees, some plots in mature stands with low tree density were excluded from the analyses, which seems to be one of the disadvantages of using small plots for fitting diameter distribution. However, due to the large total number of plots used in our study, a sufficient number of plots from mature stands (mean $DDOM$ = 42 cm, maximum $DDOM$ = 66 cm) were included in the analysis.

Several studies have confirmed that the estimation of the Weibull function parameters improves by increasing the number of trees used for fitting (Hossain & Zimmer 2003, Chu & Ke 2012, Teimouri et al. 2013, Pobočíková & Sedláčková 2014, Akram & Hayat 2014). However, this was not confirmed in our study. On the contrary, the tree density in many plots excluded from the analyses was even higher when compared to the tree density of the analyzed plots (results not shown). It seems that stand parameters affect the fitting performance of the Weibull distribution (Merganič & Sterba 2006, Nord-Larsen & Cao 2006, Schütz & Rosset 2020). Our results showed that the smallest differences be-

tween observed and predicted diameter distributions were found in stands with high BA and relatively high QMD . This means that the Weibull function fitted better to dense mature stands and worse to low density stands under regeneration that are characterized by a high QMD and a low BA (Merganič & Sterba 2006).

The irregular (e.g., bimodal or multimodal) empirical diameter distribution of trees might be the second important reason for non-convergence. Such a diameter structure may be a consequence of past disturbances, non-regular harvesting or a specific thinning regime (Merganič & Sterba 2006). Such distributions can be fitted by dividing the distribution into more parts (e.g., layer by layer) by using methods such as the finite mixture model (Teimouri & Podlaski 2022), splines or generalized additive models (Perperoglou et al. 2019) to improve fitting.

Stand heterogeneity correlates significantly with the parameter C , which indicates the shape of the diameter distribution (Bailey & Dell 1973). We found a quite weak negative correlation between the parameter C and the Gini index, indicating structural heterogeneity; the MM1 and MOM methods showed higher correlation than the MLE method. Structural diversity indices do not unambiguously identify even-aged stands; stands may be of different structures but have the same diversity index (Gove et al. 1994, Duduman 2011). Despite the low threshold value of the Gini index to identify even-aged stands (< 0.33), a surprisingly high number of plots exhibiting reversed-J-shaped distribution were found with parameter $C < 1$, especially when the MLE method was applied. One of the possible reasons is that the Gini index was calculated with the BA of trees, which means that larger trees give more weight to the index than thinner trees, while we fitted the Weibull distribution according to the number of trees. The second, but even more decisive reason for reversed-J-shaped distributions could be the presence of understorey trees (i.e., 10-20 cm in DBH), which can also be common and numerous in even-aged stands in the study area managed with the irregular shelterwood system (Klopčič & Bončina 2012). The understorey layer of small-sized trees can dramatically increase the Gini index (and structural diversity) if the threshold for tree measurement is low (O'Hara et al. 2007). Therefore, in some studies understorey trees (Kangas & Maltamo 2000) or trees below the upper canopy (Nord-Larsen & Cao 2006, Rosset et al. 2018) were excluded from the analyses, a higher diameter measurement threshold was adopted (O'Hara et al. 2007, Duduman 2011), or separate analyses were done for social classes of trees (Zasada & Cieszewski 2005).

Thirdly, the selected method for fitting the Weibull distribution function can influence non-convergence. According to Akram & Hayat (2014), the non-convergence

rate when using the MLE method is high when the plot number is small and the shape parameter $C < 1$. Our results confirmed this conclusion.

Finally, in forest inventories only trees above the DBH threshold are measured (Tomppo et al. 2008), which results in a truncated distribution of trees (Nanos & Sjöstedt De Luna 2017). To avoid bias in estimating the parameters (Zutter et al. 1986, Palahi et al. 2007), a truncated version of the (Weibull) distribution function is suggested. The bias increases as the QMD approaches the threshold measurement diameter (Zutter et al. 1986, Merganič & Sterba 2006); however, this was not the case in our study.

Conclusions

All three tested estimation methods (MM1, MOM and MLE) provided relatively reliable estimations of stand parameters (N, BA, DDOM and QMD) which are important for forest managers, but fitting the distribution to tree frequencies per DBH class exhibited much poorer results. Although MLE has been the most commonly used estimation method in forestry, its high non-convergence rate is a major disadvantage compared to the other two methods. When considering a set of criteria for the estimation of fitting methods, the MM1 method seems to be superior for fitting the three-parameter Weibull distribution to beech forests represented by data derived from small plots. Finally, data from small sample plots (0.05 ha), such as those from the NFIs, are suitable for predicting stand parameters derived from fitted or modelled diameter distributions, but less appropriate for modelling tree frequencies per DBH class.

Acknowledgements

The study was funded by the Ministry of Agriculture, Forestry and Food of Republic of Slovenia through the project V4-2014 "The development of forest models for Slovenia" (grant no. 2330-20-000234) and the Slovenian Research Agency (ARRS) program P4-0059 "Forest, forestry and renewable forest resources". The first author was partly funded by the Pahernik Foundation. A part of the study area represented the infrastructure center "Research Forest of the Department of Forestry and Renewable Forest Resources", which is part of the Network of Research and Infrastructural Centers of University of Ljubljana (MRIC UL), co-funded by the Slovenian Research Agency.

References

- Akram M, Hayat A (2014). Comparison of estimators of the Weibull distribution. *Journal of Statistical Theory and Practice* 8: 238-259. - doi: [10.1080/15598608.2014.847771](https://doi.org/10.1080/15598608.2014.847771)
- Bailey RL, Dell TR (1973). Quantifying diameter distributions with Weibull function. *Forest Sciences* 19: 97-10. [online] URL: <http://academic.oup.com/forestscience/article-abstract/19/2/97/4675411>
- Bassil S, Nyland RD, Kern CC, Kenefic LS (2019). Dynamics of the diameter distribution after selection cutting in uneven- and even-aged northern hardwood stands: a long-term evaluation. *Canadian Journal of Forest Research* 49 (12): 1525-1539. - doi: [10.1139/cjfr-2019-0204](https://doi.org/10.1139/cjfr-2019-0204)
- Bohn U, Neuhäusl R, Gollub G, Hettwer C, Neuhäuslová Z, Raus T, Schlüter H, Weber H (2000). Buchenwälder [Beech forests]. In: "Karte der natürlichen Vegetation Europas (Maßstab 1: 2.500.000)" [Map of the Natural Vegetation of Europe (scale 1:2.500.000)]. Landwirtschaftsverlag, Münster, Germany, pp. 523. [in German]
- Bončina A, Rozman A, Dakskobler I, Klopčič M, Babij V, Poljanec A (2021). Gozdni rastiščni tipi Slovenije: vegetacijske, sestojne in upravljaljske značilnosti [Forest vegetation types in Slovenia: vegetation, stand, and management characteristics]. Biotechnical Faculty, Department of Forestry and Renewable Forest Resources and Slovenia Forest Service, Ljubljana, Slovenia, pp. 575. [in Slovenian]
- Burkhardt HE, Tomé M (2012). Modeling forest trees and stands. Springer, Dordrecht, Netherlands, pp. 457. [online] URL: <http://books.google.com/books?id=XmcNcfEarsoC>
- Cao QV (2004). Predicting parameters of a Weibull function for modeling diameter distribution. *Forest Science* 50 (5): 682-685. - doi: [10.1093/forestscience/50.5.682](https://doi.org/10.1093/forestscience/50.5.682)
- Chu YK, Ke JC (2012). Computation approaches for parameter estimation of Weibull distribution. *Mathematical and Computational Applications* 17 (1): 39-47. - doi: [10.3390/mca17010039](https://doi.org/10.3390/mca17010039)
- Dubey SD (1967). Some percentile estimators for Weibull parameters. *Technometrics* 9 (1): 119-129. - doi: [10.2307/1266322](https://doi.org/10.2307/1266322)
- Duduman G (2011). A forest management planning tool to create highly diverse uneven-aged stands. *Forestry* 84 (3): 301-314. - doi: [10.1093/forestry/cpro14](https://doi.org/10.1093/forestry/cpro14)
- Gorgoso-Varela JJ, Rojo-Alboreca A, Camara-Obregon A, Dieguez-Aranda U (2012). A comparison of estimation methods for fitting Weibull, Johnson's SB and beta functions to *Pinus pinaster*, *Pinus radiata* and *Pinus sylvestris* stands in northwest Spain. *Forest Systems* 21 (3): 446-459. - doi: [10.5424/fs/2012213-02736](https://doi.org/10.5424/fs/2012213-02736)
- Gove JH, Patil GP, Swindel DF, Taillie C (1994). Ecological diversity and forest management. In: "Handbook of Statistics 12. Environmental Statistics" (Patil GP, Rao CR eds). Elsevier, Amsterdam, Netherlands, pp. 409-462. - doi: [10.1016/S0169-7161\(05\)80014-8](https://doi.org/10.1016/S0169-7161(05)80014-8)
- Hasenauer H, Kindermann G, Steinmetz P (2006). The tree growth model MOSES 3.0. In: "Sustainable Forest Management" (Hasenauer H ed). Springer, Berlin, Heidelberg, Germany, pp. 64-70. - doi: [10.1007/3-540-31304-4_5](https://doi.org/10.1007/3-540-31304-4_5)
- Hossain A, Zimmer W (2003). Comparison of estimation methods for Weibull parameters: complete and censored samples. *Journal of Statistical Computation and Simulation* 73 (2): 145-153. - doi: [10.1080/00949650215730](https://doi.org/10.1080/00949650215730)
- Kangas A, Maltamo M (2000). Performance of percentile-based diameter distribution prediction and Weibull method in independent data sets. *Silva Fennica* 24 (4): 381-398. - doi: [10.1421/4/sf.620](https://doi.org/10.1421/4/sf.620)
- Kangas A, Maltamo M (2006). Forest inventory.

- Methodology and applications. Springer, Dordrecht, Netherlands, pp. 362. - doi: [10.1007/1-4020-4381-3](https://doi.org/10.1007/1-4020-4381-3)
- Klopčič M, Bončina A (2011). Stand dynamics of silver fir (*Abies alba* Mill.)-European beech (*Fagus sylvatica* L.) forests during the past century: a decline of silver fir? *Forestry* 84 (3): 259-271. - doi: [10.1093/forestry/cpro11](https://doi.org/10.1093/forestry/cpro11)
- Klopčič M, Bončina A (2012). Recruitment of tree species in mixed selection and irregular shelterwood forest stands. *Annals of Forest Science* 69 (8): 915-925. - doi: [10.1007/s13595-012-0224-1](https://doi.org/10.1007/s13595-012-0224-1)
- Lei Y (2008). Evaluation of three methods for estimating the Weibull distribution parameters of Chinese pine (*Pinus tabulaeformis*). *Journal of Forest Science* 54 (12): 566-571. - doi: [10.17221/68/2008-JFS](https://doi.org/10.17221/68/2008-JFS)
- Merganič J, Sterba H (2006). Characterisation of diameter distribution using the Weibull function: method of moments. *European Journal of Forest Research* 125: 427-439. - doi: [10.1007/s10342-006-0138-2](https://doi.org/10.1007/s10342-006-0138-2)
- Nanos N, Sjöstedt De Luna S (2017). Fitting diameter distribution models to data from forest inventories with concentric plot design. *Forest Systems* 26 (2): 13. - doi: [10.5424/fs/2017262-10486](https://doi.org/10.5424/fs/2017262-10486)
- Nord-Larsen T, Cao VQ (2006). A diameter distribution for even-aged beech in Denmark. *Forest Ecology and Management* 231 (1-3): 218-225. - doi: [10.1016/j.foreco.2006.05.054](https://doi.org/10.1016/j.foreco.2006.05.054)
- O'Hara KL, Hasenauer H, Kindermann G (2007). Sustainability in multi-aged stands: an analysis of long-term planter systems. *Forestry* 80: 163-181. - doi: [10.1093/forestry/cpl051](https://doi.org/10.1093/forestry/cpl051)
- Palahi M, Pukkala T, Trasobares A (2006). Modelling the diameter distribution of *Pinus sylvestris*, *Pinus nigra* and *Pinus halepensis* forest stands in Catalonia using the truncated Weibull function. *Forestry* 79 (5): 553-562. - doi: [10.1093/forestry/cpl037](https://doi.org/10.1093/forestry/cpl037)
- Palahi M, Pukkala T, Blasco E, Trasobares A (2007). A comparison of beta, Johnson's SB, Weibull and truncated Weibull functions for modelling diameter distribution of forest stands in Catalonia (north-east of Spain). *European Journal of Forest Research* 126 (4): 563-571. - doi: [10.1007/s10342-007-0177-3](https://doi.org/10.1007/s10342-007-0177-3)
- Perperoglou A, Sauerbrei W, Abrahamowicz M, Schmid M (2019). A review of spline function procedures in R. *BMC Medical Research Methodology* 19: 46. - doi: [10.1186/s12874-019-0666-3](https://doi.org/10.1186/s12874-019-0666-3)
- Pobočková I, Sedláčková Z (2014). Comparison of four methods for estimating the Weibull distribution parameters. *Applied Mathematical Sciences* 8 (83): 4137-4149. - doi: [10.12988/ams.2014.45389](https://doi.org/10.12988/ams.2014.45389)
- Poudel KP, Cao QV (2013). Evaluation of methods to predict Weibull parameters for characterizing diameter distributions. *Forest Science* 59 (2): 243-252. - doi: [10.5849/forsci.12-001](https://doi.org/10.5849/forsci.12-001)
- Pretzsch H, Biber P, Dursky J (2002). The single tree-based stand simulator SILVA: construction, application and evaluation. *Forest Ecology and Management* 162 (1): 3-21. - doi: [10.1016/S0378-1127\(02\)00047-6](https://doi.org/10.1016/S0378-1127(02)00047-6)
- Pretzsch H (2009). Forest dynamics, growth and yield. From measurement to model. Springer, Berlin, Heidelberg, Germany, pp. 664. - doi: [10.1007/978-3-540-88307-4](https://doi.org/10.1007/978-3-540-88307-4)
- Rosset C, Dumollard G, Endtner J, Gollub C, Mar-

- tin V, Sala Weber V D, Wyss F, Schütz JP (2018). SiWaWa 2.0 et Placettes Permanentes de Suivi Sylvicole (PPSS). Rapport Final, SiWaWa, un modèle d'inventaire et de croissance à l'échelle du peuplement 2018 [SiWaWa 2.0 and Permanent Silvicultural Monitoring Plots (PPSS). Final Report, SiWaWa, a stand-scale inventory and growth model 2018]. Office fédéral de l'environnement - OFEV, Division Forêts, Berne, Switzerland, pp. 127. [in French]
- Schmidt LN, Sanquetta MNI, McTague JP, Silva GF, Fraga Filho CV, Sanquetta CR, Soares Scolforo JR (2020). On the use of the Weibull distribution in modeling and describing diameter distributions of clonal eucalypt stands. *Canadian Journal of Forest Research* 50 (10): 1050-1063. - doi: [10.1139/cjfr-2020-0051](https://doi.org/10.1139/cjfr-2020-0051)
- Schütz JP, Rosset C (2020). Performances of different methods of estimating the diameter distribution based on simple stand structure variables in monospecific regular temperate European forests. *Annals of Forest Science* 77, 47. - doi: [10.1007/s13595-020-00951-3](https://doi.org/10.1007/s13595-020-00951-3)
- Senthamarai Kannan K, Manoj K, Arumugam S (2015) Labeling Methods for Identifying Outliers. *International Journal of Statistics and Systems* 10 (2): 231-238. [online] URL: <https://www.researchgate.net/publication/283755180>
- SFS (2021). Letno poročilo o gozdovih za leto 2020 [Annual report on forests for 2020]. Slovenia Forest Service, Ljubljana, Slovenia, pp. 125. [in Slovenian]
- Sghaier T, Cañellas I, Calama Sánchez-González M (2016). Modelling diameter distribution of *Tetraclinis articulata* in Tunisia using normal and Weibull distributions with parameters depending on stand variables. *iForest* 9: 702-709. - doi: [10.3832/for1688-008](https://doi.org/10.3832/for1688-008)
- Siipilehto J (1999). Improving the accuracy of predicted basal-area diameter distribution in advanced stands by determining stem number. *Silva Fennica* 33 (4): 281-301. - doi: [10.14214/sf.650](https://doi.org/10.14214/sf.650)
- Sterba H, Moser M, Hasenauer H, Monserud RA (1995). PROGNAUS ein abstandsunabhängiger Wachstumssimulator für ungleichaltrige Mischbestände [PROGNAUS a distance-independent growth simulator for mixed stands of the same age]. Deutscher Verband Forstlicher Forschungsanstalten, Sektion Ertragskunde, Eberswald, Berlin, Germany, pp. 173-183. [in German]
- Teimouri M, Seyed M, Hoseini SM, Nadarajah S (2013). Comparison of estimation methods for the Weibull distribution statistics. *Journal of Theoretical and Applied Statistics* 47 (1): 93-109. - doi: [10.1080/02331888.2011.559657](https://doi.org/10.1080/02331888.2011.559657)
- Teimouri M (2020). ForestFit: statistical modeling for plant size distributions. R package version 1.2.3, web site. [online] URL: <http://cran.r-project.org/package=ForestFit>
- Teimouri M, Podlaski R (2022). Bayesian Inference for Johnson's SB and Weibull distributions. *Scandinavian Journal of Forest Research* 37 (1): 74-82. - doi: [10.1080/02827581.2021.2005132](https://doi.org/10.1080/02827581.2021.2005132)
- Tomppo E, Haakana M, Katila M, Peräsaari J (2008). Multi-source national forest inventory - Methods and applications. Springer, Dordrecht, Netherlands, pp. 373. - doi: [10.1007/978-1-4020-8713-4](https://doi.org/10.1007/978-1-4020-8713-4)
- Weibull W (1939). A statistical theory of the strength of material. *Ingeniors Vetenskaps Akademiens Handlingar* 151: 1-45.
- Zanakis SH (1979). A simulation study of some simple estimators for the three-parameter Weibull distribution. *Journal of Statistical Computation and Simulation* 9: 101-116. - doi: [10.1080/00949657908810302](https://doi.org/10.1080/00949657908810302)
- Zasada M, Cieszewski CJ (2005). A finite mixture distribution approach for characterizing tree diameter distributions by natural social class in pure even-aged Scots pine stands in Poland. *Forest Ecology and Management* 204 (2-3): 145-158. - doi: [10.1016/j.foreco.2003.12.023](https://doi.org/10.1016/j.foreco.2003.12.023)
- Zhang L, Packard KC, Liu C (2003). A comparison of estimation methods for fitting Weibull and Johnson's SB distributions to mixed spruce-fir stands in northeastern North America. *Canadian Journal of Forest Research* 33: 1340-1347. - doi: [10.1139/X03-054](https://doi.org/10.1139/X03-054)
- Zutter BR, Oderwald RG, Murphy PA, Farrar RM (1986). Characterizing diameter distributions with modified data types and forms of the Weibull distribution. *Forest Science* 32 (1): 37-48. - doi: [10.1093/forestscience/32.1.37](https://doi.org/10.1093/forestscience/32.1.37)
- Zuur AF, Ieno EN, Smith GM (2007). *Analysing ecological data*. Springer, New York, USA, pp. 672. - doi: [10.1007/978-0-387-45972-1](https://doi.org/10.1007/978-0-387-45972-1)