

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/320740765>

# Hydration, drinking and exercise performance

Article · January 2017

---

CITATION  
1

READS  
218

source: <https://doi.org/10.24451/arbor.6998> | downloaded: 18.7.2024

# Hydration, drinking and exercise performance

Mettler S<sup>1,2</sup>, Mannhart Ch<sup>3</sup>

<sup>1</sup> Department of Business, Health and Social Work, Bern University of applied sciences, Switzerland, samuel.mettler@bfh.ch;

<sup>2</sup> Department of Health Sciences and Technology, ETH Zurich, Switzerland, samuel-mettler@ethz.ch;

<sup>3</sup> consultingmannhart, Wolfhausen, Switzerland, mail@consultingmannhart.ch

## Abstract

To counter progressive dehydration and thirst, athletes drink during exercise. However, despite decades of scientific research, there is still no conclusive answer regarding how much we should drink to optimize performance. The goal of this review article is to analyze the arguments underpinning contrasting perspectives and to critically analyze the available evidence.

It seems that the respective argumentations of contrasting viewpoints are based on a different selective fraction of the available evidence. In studies using time trial performance protocols in which dehydration develops during exercise, it seems that end-exercise dehydration levels of up to 4% body mass do not compromise endurance performance in temperate to hot conditions – at least as long as the athlete is not prevented from drinking. In contrast, studies that induced dehydration pre-exercise consistently report performance impacts already at low levels of dehydration, i.e., 1 to 2% body mass loss.

Further factors like the perception of thirst have been suggested to influence performance, but performance effects cannot be explained solely by the perception of thirst as well. Nevertheless, no evidence was found against the hypothesis that drinking ad libitum may optimize performance outcomes. At the same time, arguments have been identified regarding why a drinking plan might assist athletes in different situations.

## Zusammenfassung

Athletinnen und Athleten trinken während dem Sport, um Flüssigkeitsdefizite und Durst zu reduzieren oder zu vermeiden. Trotz jahrzehntelanger Forschung gibt es nach wie vor unterschiedliche Sichtweisen darüber, wie viel während dem Sport getrunken werden soll, um die Leistung zu optimieren. Das Ziel dieses Reviewartikels besteht darin, die Argumente hinter sich widersprechenden Sichtweisen zu beleuchten und die verfügbare Evidenz kritisch zu hinterfragen.

Sich widersprechende Theorien scheinen auf einer unterschiedlichen selektiven Auswahl aus der verfügbaren Evidenzgrundlage zu basieren. In Studien, welche die Leistung mit Time Trials gemessen haben und sich das Flüssigkeitsdefizit während Belastung entwickelte, konnte bei milden und warmen Umgebungsbedingungen bis zu einem Flüssigkeitsdefizit von 4% der Körpermasse kein negativer Effekt auf die Ausdauerleistungsfähigkeit festgestellt werden. Dies zumindest so lange, als die Athleten nicht am Trinken gehindert werden. Im Gegensatz dazu wurde in Studien, welche das Flüssigkeitsdefizit vor Belastungsstart induziert haben, konsistent bereits bei Flüssigkeitsdefiziten von 1 bis 2% der Körpermasse eine Leistungsbeeinträchtigung festgestellt.

Weitere Faktoren wie die Durstwahrnehmung könnten die Leistung mit beeinflussen, wobei Leistungseffekte aber auch nicht alleine aufgrund des Durstgefühls erklärt werden können. Allerdings konnten keine Hinweise gegen die Hypothese gefunden werden, wonach gemäss Durst zu trinken die Leistung optimieren würde. Gleichzeitig gibt es auch Argumente, welche für die Ausarbeitung eines Trinkplans sprechen.

## Introduction

Sweat evaporation provides an important heat dissipation avenue, particularly during high exercise intensity [1]. Prolonged sweating may result in increasing levels of dehydration, which has been associated with effects on both exercise performance and thermoregulation [1]. To counter progressive dehydration, athletes voluntarily drink as a response to their perception of thirst and because of externally applied drinking guidelines. The goal of this review article is to critically analyze the available evidence about the influence of hydration and drinking on exercise performance. Fueling effects by ingesting carbohydrates together with the fluid are not in the focus of this article.

## The history of drinking

Until the late 1960s athletes were convinced that avoiding drinking during exercise would optimize performance [2]. In 1975 by the American College of Sport Medicine (ACSM) published its first drinking guidelines. Those guidelines focused on the avoidance of “heat injuries” [3]. Performance was not in the focus until the ACSM’s 1996 position stand [4]. Notably, the 1996 guidelines suggested that “individuals should be encouraged to consume the maximal amount of fluids during exercise that can be tolerated without gastrointestinal discomfort up to a rate equal to that lost from sweating.” Absolute drinking volumes of 0.6 to 1.2 liters per hour of exercise were recommended.

These drinking guidelines have been seriously questioned [5–8]. In addition to the criticisms of the scientific evidence for these guidelines, critics have highlighted the danger of overhydration and hyponatremia [7,9]. Indeed, at the beginning of the new millennium, overhydration and associated hyponatremia have become a mass phenomenon [10–13], indicating that the advice to drink during exercise has obviously taken root in the population around the world, but perhaps not as intended by the authors of the message. Field studies in endurance events around the world started to report a significant proportion of participants finishing the events overhydrated [11–14], with an increasing risk for asymptomatic and symptomatic hyponatremia and even fatal outcomes [13,15,16].

The 2007 ACSM position stand on exercise and fluid replacement [1] suggested ad libitum drinking from 0.4 to 0.8 liters per hour for endurance events. It was suggested to keep body mass loss <2% in order to avoid performance decrements. These arguments were confirmed in the latest ACSM position stand [17] and represent the classical perspective on the topic of dehydration, drinking and performance.

## Two fundamental perspectives

Since the 1990s the classical perspective has been more and more challenged. After nearly a century of scientific research, numerous studies have been published about the influence of fluid intake and dehydration on diverse exercise performance and thermoregulatory outcomes [18–86]. Review articles and meta-analyses have tried to compile the data and to draw practical conclusions, but actually ended up with conflicting interpretations [87–96] and the scientific community is far from a common position. Instead, two fun-

### The classical perspective

The classical view, as represented by both the 2007 [1] and 2016 [16] ACSM position stand and additional review articles about the influence of drinking or dehydration on thermoregulation and performance [86,87,93–95] adopt the following fundamental line of arguments:

1. Fluid deficits of more than 1–2% body mass impair exercise performance, in particular endurance exercise and under warm conditions.
2. Fluid deficits increase body core temperatures and thereby increase cardiovascular stress, which may impair performance and increase the risk for heat illness or heat stroke.
3. Drinking “ad libitum” or “to thirst” is not sufficient to prevent heat illness or performance decrements.

### The contrasting perspective

The contrasting view challenges the classical view and provides the following line of arguments [2,6,7,13,86,88–90,92]:

1. Fluid deficits of more than 2% body mass do not necessarily impair exercise performance.
2. There is no evidence that drinking more than ad libitum or to thirst provides any performance advantage over paced fluid ingestion to minimize body mass loss to 1–2%.
3. Exercise intensity, not dehydration, is the primary cause for heat stroke.

**Table 1:** Two perspectives on the effect of drinking and dehydration on exercise performance and thermoregulation.

damentally different viewpoints (*see Tab. 1*) have emerged, which have not significantly changed over the last decade [87–89].

## Evidence for real-life endurance performance

To allow for transferring conclusions from laboratory-based research to real-life settings, the study design should simulate real life as much as possible (i.e., the external or ecologic validity of the study). One difference between many laboratory studies and real-life settings is that the majority of studies [20–23,27–31,33,35,38–40,42–45,47–54,56–58,61,62,66,67,69–71,73,77,82,84,86] induced dehydration before exercise, either by exercising the subject in the heat, imposing fluid restrictions of about one to two days, or administering diuretic drugs before the exercise task. This is in contrast to real life, where athletes dehydrate during exercise (exercise-induced dehydration). Further, in real life, athletes have to perform as fast as possible over a given distance while being able to self-pace their race and allow continual behavioral adjustments. In contrast, many laboratory studies used fixed-power output tests to exhaustion, which only allow an all-or-none response (exercise continuation or termination). There is no real-world sport where athletes need to perform as long as possible without being able to adjust power output and without knowing the distance or time (workload) to be covered. The ecologic validity of time to exhaustion tests, particularly combined with pre-exercise dehydration, is therefore considered as low [90,93,97].

When screening the literature for studies using time trial performance and exercise-induced dehydration, not many studies remain [19,37,60,63–65,75,76,79–81,83]. Interestingly, these studies reported no performance impact of dehydration with body mass losses up to 2% [76,79,81,83], beyond 2% [37,63,65,80], and up to 3.2% [41,60,75] or 4% [64], covering temperate to hot conditions. Exceptions are Below et al.'s [19] and Dugas et al.'s [80] studies, where drinking nothing or below ad libitum was associated with reduced performance.

Taken together, these results suggest that under lab conditions that consider two important real-life settings, end-exercise dehydration levels of up to 4% body mass do not seem to compromise endurance performance in temperate to hot conditions, at least as long as drinking is not withheld from the athlete. Two recent meta-analyses come to a similar conclusion, suggesting that studies with ecologically valid study designs do not show performance impacts of exercise-induced dehydration of up to 4% body mass [90,91], whereas studies using protocols with fixed exercise intensity and/or pre-exercise induced dehydration, 1–2% body mass loss is already associated with reduced performance [39,40,91].

Observational field data of elite endurance athletes winning international marathons have reported dehydration levels of 6.6% to 11.7% body mass [98]. Although these observations do not allow for a causative conclusion that dehydration does not influence performance, it is nevertheless a striking contrast to the classical viewpoint, which suggests that endurance performance would be impaired if dehydration exceeds 1–2% body mass loss [1].

In the 2007 ACSM position stand [1], three reviews [95,99,100] were cited for the conclusion that dehydration of >2% body mass degrades aerobic exercise performance in warm environments. The cited reviews themselves refer to further reviews and studies based on ecologically invalid study designs (e.g. time to exhaustion tests and or induction of dehydration by artificial pre-exercise protocols).

One interpretation of the conflicting viewpoints (*Tab. 1*) is that they are simply founded on different study types and a different body of evidence, which separate themselves by their ecologic validity.

## Dehydration, thirst and performance

Dugas et al. [80] showed that two hour cycling time trial performance is optimized when drinking ad libitum or more during exercise, compared with restricted fluid (no fluid or less than ad libitum). Noakes [7,8] has suggested that the optimal amount of drinking is ad libitum or to thirst. In contrast, Backx et al. [81] found no effect of drinking more or less than ad libitum during a one hour cycling time trial. Daries et al. [75] or Dion et al. [60] found no positive performance effect on a two hour cycling time trial [75] or 21 km running [60] when ad libitum drinking of about 0.4 l/h was forced to 0.9 l/h or 1.3 l/h, respectively, in order to reduce end-exercise dehydration. Nevertheless, all three studies are in line with the observation that no single intervention study has ever been published showing that drinking more than ad libitum or to thirst would be associated with any performance benefits, irrespective of the end-exercise dehydration level. On the other hand, drinking less than ad libitum may [19,80] or may not [63,76,79,81,83] negatively affect performance in ecologically valid studies.

Interestingly, the perception of thirst may be reduced with increased drinking, but without affecting performance [60,64]. Simultaneously, increasing abdominal discomfort may occur [60]. This implies that maximal reduction of thirst is not the chosen level of the thirst signal when drinking ad libitum. However, it may represent the optimal drinking level from an integrated perspective of different body signals.

Fallowfield et al. [24] let subjects run to exhaustion at 70%  $\text{VO}_2\text{max}$  either by drinking nothing or by drinking a moderate amount (0.5 l/h) of water. With drinking, subjects ran 34% longer (103 vs. 77 min). Due to the longer running time, the resulting end-exercise dehydration was increased compared to the non-drinking trial. This time to exhaustion study supports the hypothesis that the dehydration level is an inferior parameter compared to ingesting fluids. Unfortunately, thirst was not assessed in this study.

In conclusion, it seems that neither dehydration nor thirst can explain performance effects on their own in laboratory-based intervention studies. Further studies with exercise durations of at least 2 h and ecologically valid study designs are needed to further understand the interaction of these parameters.

## Challenges of drinking ad libitum in the field

From a laboratory-based perspective, drinking ad libitum seems to be an optimal drinking strategy. However, it has been questioned whether performance is optimized just by drinking ad libitum in all sports and in all individual real-life situations, as e.g., game rules or tactical constraints may restrict drinking to certain time points [101]. An alternative interpretation is that the definition of the term “ad libitum” is broadened by the fact that both internal body signals and external constraints are integrated to an optimal behavior.

In contrast to the two fundamental viewpoints (*Tab. 1*), we do not see an irreconcilable incompatibility between drinking ad libitum and establishing an individualized drinking plan. Rather, without a plan we see some critical points that may arise in real life compared to laboratory-based endurance tests. We could also imagine that some of these points may interfere with the brain's ability to optimally integrate all signals, including thirst sensation, into a suitable drinking behavior. Finally, ad libitum drinking is not completely random and may become reasonably predictable after acquiring data about the athletes' drinking and body mass (see e.g. drinking calculator on <http://www.sns.ch/sportsnutrition/trinkmengenrechner>). Transferring this experience into a plan may allow for planning and organization, avoiding distraction from drinking, ensuring optimal fueling, and providing guidance to athletes who are nervous or inexperienced with the exercise task to come. At the available time points, drinking may still be ad libitum. There are also real-life situations in which there may be significant arguments to drink beyond ad libitum. I.e., in tournament or multiple training situation quick recovery is easier when avoiding exaggerated dehydration, which might not be a performance problem for a single exercise task.

## Tolerable upper drinking levels

Gastric emptying rates depend on, e.g., gastric filling [102], the carbohydrate content of the drink [103], and solute osmo-

lality [104]. With repeated drinking, probably around 1.2 l/h may pass the stomach with an acceptable stomach filling during moderate intensive exercise [104].

Laboratory based studies reported gastrointestinal problems or feelings of fullness for drinking 0.9 l/h [75,81] or more [37,60,105] during running or cycling time-trials. Reported ad libitum fluid intake in laboratory based studies was between 0.4 and 0.7 l/h for one to two hours high intensity endurance running, cycling or soccer [60,63,75,80,81]. This corresponds with field observations reporting fluid intakes from 0.2 l/h to 0.8 l/h for various exercise [101]. Compared to running, higher fluid intake rates of around 1.0 l/h are occasionally observed on the bike [101], and triathletes drink more during the bike stage than during running [106]. With frequent drinking during regular recovery breaks intermittent sports seem to allow fluid intake rates of beyond 1 l/h (e.g., tennis or basketball) [101,107].

### Dehydration and performance in non-endurance sports

Pre-exercise induced dehydration of  $\geq 2\%$  body mass resulted in consistent reports of reduced basketball skill performance and sprint times [49,51], riding [62] or golf-specific motor and mental performance [66]. Studies about basketball- and soccer-specific exercise tasks and skills while applying during-exercise dehydration are less consistent [46,55,63,74,85], with some finding no effect on the Loughborough intermittent shuttle running test (LIST) [63,85], while others reported negative effects of drinking nothing and concomitant increased dehydration on goals shot and anaerobic or sprint performance [46,55]. The difficulty of interpreting these studies is that they compared drinking nothing at all versus drinking mostly equivalent to 100% of the sweat losses. Therefore, the only practical conclusion is that drinking nothing is probably not ideal. The only study testing ad libitum drinking vs. nothing vs. equivalent to sweat losses found no difference on LIST performance for any drinking schedule [63].

A comprehensive meta-analysis of the influence of dehydration on muscle endurance, strength, anaerobic performance, and jump performance reported negative effects of dehydration of  $>2\%$  body mass on muscle endurance, strength, and anaerobic performance [92], with some earlier reviews coming to similar conclusions [52,108]. In contrast, vertical jump ability was non-significantly improved with an average dehydration of 2.7 % body mass [92]. Regarding anaerobic sprint performance, it is worth noting that single sprints [45,47,48] seem to be rather unaffected by dehydration compared to repeated sprints or intermittent exercise tasks [44,50,54,68].

However, the ecologic validity of all studies mentioned above or included in Savoie et al.'s [92] meta-analysis may be questioned, as dehydration was induced by pre-exercise dehydration protocols involving heat exposure with or without endurance exercise tasks or fluid restriction of up to three days.

During-exercise dehydration is more difficult to induce in short, explosive exercise tasks compared to endurance exercise. Nevertheless, real-life competition settings may still last several hours, including preparation, sport-specific warm-up procedures, and competition with, e.g., six attempts in long jump. During these procedures, significant dehydration may undoubtedly occur and pronounced dehydration levels may be seen especially in combined events (personal observa-

tions). Regrettably, if a real-life athlete, e.g., a long jumper, would like to know whether dehydration influences his performance, we probably have to conclude that after decades of modern research, the available evidence does not allow for a conclusive answer.

### Conclusion and practical implications

Based on the currently available evidence, one can conclude that drinking ad libitum is the best general advice to give to athletes. Absolute drinking volumes are difficult to recommend, although typical drinking volumes are often in the range of 0.4 to 0.8 l/h. Nevertheless, ad libitum or required drinking volumes may be below or beyond this range in some situations, and in contrast to ACSM's line of arguments, substantial evidence has emerged that exercise performance is not necessarily compromised when reaching dehydration levels of beyond 2% body mass. Further, drinking ad libitum and establishing an individualized drinking plan are compatible. Indeed, ad libitum drinking may be optimized with a drinking plan in real-life and there are situations where drinking beyond ad libitum might be advisable.

### Corresponding author

Samuel Mettler  
Department of Business, Health and Social Work  
Bern University of applied sciences  
E-Mail: samuel.mettler@bfh.ch

Department of Health Sciences and Technology  
ETH Zurich  
E-Mail: samuel-mettler@ethz.ch

### References

1. Sawka MN, Burke LM, Eichner ER, Maughan RJ, Montain SJ, and Stachenfeld NS. Exercise and fluid replacement. *Med Sci Sports Exerc.* 2007;39:377-390.
2. Noakes TD. Fluid replacement during exercise. *Exerc Sport Sci Rev.* 1993;21:297-330.
3. Medicine ACoS. The american college of sports medicine position statement on prevention of heat injuries during distance running. *Med Sci Sports.* 1975;7:Vii-ix.
4. Convertino VA, Armstrong LE, Coyle EF, Mack GW, Sawka MN, Senay LC, Jr., and Sherman WM. American college of sports medicine position stand. Exercise and fluid replacement. *Med Sci Sports Exerc.* 1996;28:i-vii.
5. Noakes TD. Dehydration during exercise: What are the real dangers? *Clin J Sport Med.* 1995;5:123-8.
6. Pitsiladis Y and Beis L. To drink or not to drink to drink recommendations: The evidence. *Brit Med J.* 2012;345:e4868.
7. Noakes TD. Is drinking to thirst optimum? *Ann Nutr Metab* 2010;57 Suppl 2:9-17.
8. Noakes TD. Drinking guidelines for exercise: What evidence is there that athletes should drink "as much as tolerable", "to replace the weight lost during exercise" or "ad libitum"? *J Sports Sci.* 2007;25:781-796.
9. Noakes TD and Speedy DB. Case proven: Exercise associated hyponatraemia is due to overdrinking. So why did it take 20 years before the original evidence was accepted? *Brit J Sports Med.* 2006;40:567-572.
10. Hew-Butler T, Ayus JC, Kipps C, Maughan RJ, Mettler S, Meeuwisse WH, Page AJ, Reid SA, Rehrer NJ, Roberts WO, Rogers IR, Rosner MH, Siegel AJ, Speedy DB, Stuempfle KJ, Verbalis JG, Weschler LB, and Wharam P. Statement of the second international exercise-associated hyponatremia consensus development conference, new zealand, 2007. *Clin J Sport Med.* 2008;18:111-121.

11. Mettler S, Rusch C, Frey WO, Bestmann L, Wenk C, and Colombani PC. Hyponatremia among runners in the zurich marathon. *Clin J Sport Med.* 2008;18:344-349.
12. Almond CS, Shin AY, Fortescue EB, Mannix RC, Wypij D, Binstadt BA, Duncan CN, Olson DP, Salerno AE, Newburger JW, and Greenes DS. Hyponatremia among runners in the boston marathon. *N Engl J Med.* 2005;352:1550-1556.
13. Noakes TD. *Waterlogged.* United States of America: Human Kinetics; 2012
14. Noakes TD, Sharwood K, Speedy D, Hew T, Reid S, Dugas J, Almond C, Wharam P, and Weschler L. Three independent biological mechanisms cause exercise-associated hyponatremia: Evidence from 2,135 weighed competitive athletic performances. *Proc Natl Acad Sci U S A.* 2005;102:18550-18555.
15. Kipps C, Sharma S, and Tunstall PD. The incidence of exercise-associated hyponatraemia in the london marathon. *Br J Sports Med.* 2009;Epub.
16. Hew-Butler T, Rosner MH, Fowkes-Godek S, Dugas JP, Hoffman MD, Lewis DP, Maughan RJ, Miller KC, Montain SJ, Rehrer NJ, Roberts WO, Rogers IR, Siegel AJ, Stuempfle KJ, Winger JM, and Verbalis JG. Statement of the 3rd international exercise-associated hyponatremia consensus development conference, carlsbad, california, 2015. *Br J Sports Med.* 2015;49:1432-46.
17. Thomas DT, Erdman KA, and Burke LM. Position of the academy of nutrition and dietetics, dietitians of canada, and the american college of sports medicine: Nutrition and athletic performance. *J Acad Nutr Diet.* 2016;116:501-28.
18. Barr SI, Costill DL, and Fink WJ. Fluid replacement during prolonged exercise: Effects of water, saline, or no fluid. *Med Sci Sports Exerc.* 1991;23:811-7.
19. Below PR, Mora-Rodriguez R, Gonzalez-Alonso J, and Coyle EF. Fluid and carbohydrate ingestion independently improve performance during 1 h of intense exercise. *Med Sci Sports Exerc.* 1995;27:200-210.
20. Burge CM, Carey MF, and Payne WR. Rowing performance, fluid balance, and metabolic function following dehydration and rehydration. *Med Sci Sports Exerc.* 1993;25:1358-64.
21. Buskirk ER, Iampietro PF, and Bass DE. Work performance after dehydration: Effects of physical conditioning and heat acclimatization. *J Appl Physiol.* 1958;12:189-94.
22. Caldwell JE, Ahonen E, and Nousiainen U. Differential effects of sauna-, diuretic-, and exercise-induced hypohydration. *J Appl Physiol Respir Environ Exerc Physiol.* 1984;57:1018-23.
23. Chevront SN, Carter R, 3rd, Castellani JW, and Sawka MN. Hypohydration impairs endurance exercise performance in temperate but not cold air. *J Appl Physiol.* 2005;99:1972-6.
24. Fallowfield JL, Williams C, Booth J, Choo BH, and Grows S. Effect of water ingestion on endurance capacity during prolonged running. *J Sports Sci.* 1996;14:497-502.
25. McConell GK, Burge CM, Skinner SL, and Hargreaves M. Influence of ingested fluid volume on physiological responses during prolonged exercise. *Acta Physiol Scand.* 1997;160:149-156.
26. Montain SJ and Coyle EF. Influence of graded dehydration on hyperthermia and cardiovascular drift during exercise. *J Appl Physiol.* 1992;73:1340-1350.
27. Montain SJ, Smith SA, Mattot RP, Zientara GP, Jolesz FA, and Sawka MN. Hypohydration effects on skeletal muscle performance and metabolism: A 31p-mrs study. *J Appl Physiol.* 1998;84:1889-94.
28. Saltin B. Aerobic and anaerobic work capacity after dehydration. *J Appl Physiol.* 1964;19:1114-8.
29. Slater GJ, Rice AJ, Sharpe K, Tanner R, Jenkins D, Gore CJ, and Hahn AG. Impact of acute weight loss and/or thermal stress on rowing ergometer performance. *Med Sci Sports Exerc.* 2005;37:1387-94.
30. Slater GJ, Rice AJ, Tanner R, Sharpe K, Jenkins D, and Hahn AG. Impact of two different body mass management strategies on repeat rowing performance. *Med Sci Sports Exerc.* 2006;38:138-146.
31. Craig EN and Cummings EG. Dehydration and muscular work. *J Appl Physiol.* 1966;21:670-674.
32. Ladell WS. The effects of water and salt intake upon the performance of men working in hot and humid environments. *J Physiol.* 1955;127:11-46.
33. Nybo L, Jensen T, Nielsen B, and Gonzalez-Alonso J. Effects of marked hyperthermia with and without dehydration on vo(2) kinetics during intense exercise. *J Appl Physiol.* 2001;90:1057-64.
34. Gonzalez-Alonso J, Mora-Rodriguez R, Below PR, and Coyle EF. Dehydration markedly impairs cardiovascular function in hyperthermic endurance athletes during exercise. *J Appl Physiol (1985).* 1997;82:1229-36.
35. Pichan G, Gauttam RK, Tomar OS, and Bajaj AC. Effect of primary hypohydration on physical work capacity. *Int J Biometeorol.* 1988;32:176-180.
36. Walsh RM, Noakes TD, Hawley JA, and Dennis SC. Impaired high-intensity cycling performance time at low levels of dehydration. *Int J Sports Med.* 1994;15:392-398.
37. Robinson TA, Hawley JA, Palmer GS, Wilson GR, Gray DA, Noakes TD, and Dennis SC. Water ingestion does not improve 1-h cycling performance in moderate ambient temperatures. *Eur J Appl Physiol Occup Physiol.* 1995;71:153-60.
38. Armstrong LE, Costill DL, and Fink WJ. Influence of diuretic-induced dehydration on competitive running performance. *Med Sci Sports Exerc.* 1985;17:456-61.
39. Bardis CN, Kavouras SA, Kostis L, Markousi M, and Sidossis LS. Mild hypohydration decreases cycling performance in the heat. *Med Sci Sports Exerc.* 2013;45:1782-9.
40. Bardis CN, Kavouras SA, Arnaoutis G, Panagiotakos DB, and Sidossis LS. Mild dehydration and cycling performance during 5-kilometer hill climbing. *J Athl Train.* 2013;48:741-7.
41. Cheung SS, McGarr GW, Mallette MM, Wallace PJ, Watson CL, Kim IM, and Greenway MJ. Separate and combined effects of dehydration and thirst sensation on exercise performance in the heat. *Scandinavian Journal of Medicine & Science in Sports.* 2015;25 Suppl 1:104-11.
42. Ribisl PM and Herbert WG. Effects of rapid weight reduction and subsequent rehydration upon the physical working capacity of wrestlers. *Res Q.* 1970;41:536-41.
43. Nielsen B, Kubica R, Bonnesen A, Rasmussen IB, Stoklosa J, and Wilk B. Physical work capacity after dehydration and hyperthermia. *Scand J Sports Sci.* 1981;3:2-10.
44. Davis JK, Laurent CM, Allen KE, Green JM, Stolworthy NI, Welch TR, and Nevett ME. Influence of dehydration on intermittent sprint performance. *J Strength Cond Res.* 2015;29:2586-93.
45. Chevront SN, Carter R, III, Haymes EM, and Sawka MN. No effect of moderate hypohydration or hyperthermia on anaerobic exercise performance. *Med Sci Sports Exerc.* 2006;38:1093-1097.
46. Hoffman JR, Stavsky H, and Falk B. The effect of water restriction on anaerobic power and vertical jumping height in basketball players. *Int J Sports Med.* 1995;16:214-8.
47. Jacobs I. The effects of thermal dehydration on performance of the wingate anaerobic test. *Int J Sports Med.* 1980;1:21-24.
48. Watson G, Judelson DA, Armstrong LE, Yeargin SW, Casa DJ, and Maresh CM. Influence of diuretic-induced dehydration on competitive sprint and power performance. *Med Sci Sports Exerc.* 2005;37:1168-74.
49. Baker LB, Dougherty KA, Chow M, and Kenney WL. Progressive dehydration causes a progressive decline in basketball skill performance. *Med Sci Sports Exerc.* 2007;39:1114-1123.
50. Kraft JA, Green JM, Bishop PA, Richardson MT, Neggens YH, and Leeper JD. Effects of heat exposure and 3% dehydration achieved via hot water immersion on repeated cycle sprint performance. *J Strength Cond Res.* 2011;25:778-86.
51. Dougherty KA, Baker LB, Chow M, and Kenney WL. Two percent dehydration impairs and six percent carbohydrate drink improves boys basketball skills. *Med Sci Sports Exerc.* 2006;38:1650-1658.
52. Judelson DA, Maresh CM, Anderson JM, Armstrong LE, Casa DJ, Kraemer WJ, and Volek JS. Hydration and muscular performance: Does fluid balance affect strength, power and high-intensity endurance? *Sports Med.* 2007;37:907-921.
53. Kraft JA, Green JM, Bishop PA, Richardson MT, Neggens YH, and Leeper JD. Impact of dehydration on a full body resistance exercise protocol. *Eur J Appl Physiol.* 2010;109:259-67.
54. Maxwell NS, Gardner F, and Nimmo MA. Intermittent running: Muscle metabolism in the heat and effect of hypohydration. *Med Sci Sports Exerc.* 1999;31:675-83.
55. Mohr M, Mujika I, Santisteban J, Randers MB, Bischoff R, Solano R, Hewitt A, Zubillaga A, Peltola E, and Krstrup P. Examination of fatigue development in elite soccer in a hot environment: A multi-experimental approach. *Scandinavian Journal of Medicine & Science in Sports.* 2010;20 Suppl 3:125-32.
56. Stewart CJ, Whyte DG, Cannon J, Wickham J, and Marino FE. Exercise-induced dehydration does not alter time trial or neuromuscular performance. *Int J Sports Med.* 2014;35:725-30.
57. Fleming J and James LJ. Repeated familiarisation with hypohydration attenuates the performance decrement caused by hypohydration during treadmill running. *Appl Physiol Nutr Metab.* 2014;39:124-9.
58. Wilk B, Meyer F, Bar-Or O, and Timmons BW. Mild to moderate hypohydration reduces boys' high-intensity cycling performance in the heat. *Eur J Appl Physiol.* 2014;114:707-13.
59. Ramos-Jimenez A, Hernandez-Torres RP, Wall-Medrano A, Torres-Duran PV, Juarez-Oropeza MA, and Solis Ceballos JA. Acute physiological response to indoor cycling with and without hydration; case and self-control study. *Nutr Hosp.* 2013;28:1487-93.

60. Dion T, Savoie FA, Asselin A, Garipey C, and Goulet ED. Half-marathon running performance is not improved by a rate of fluid intake above that dictated by thirst sensation in trained distance runners. *Eur J Appl Physiol*. 2013;113:3011-20.
61. Wall BA, Watson G, Peiffer JJ, Abbiss CR, Siegel R, and Laursen PB. Current hydration guidelines are erroneous: Dehydration does not impair exercise performance in the heat. *Brit J Sports Med*. 2015;49:1077-83.
62. Wilson G, Hawken MB, Poole I, Sparks A, Bennett S, Drust B, Morton J, and Close GL. Rapid weight-loss impairs simulated riding performance and strength in jockeys: Implications for making-weight. *J Sports Sci*. 2014;32:383-91.
63. Owen JA, Kehoe SJ, and Oliver SJ. Influence of fluid intake on soccer performance in a temperate environment. *J Sports Sci*. 2013;31:1-10.
64. Munoz CX, Carney KR, Schick MK, Coburn JW, Becker AJ, and Judelson DA. Effects of oral rehydration and external cooling on physiology, perception, and performance in hot, dry climates. *Scandinavian Journal of Medicine & Science in Sports*. 2012;22:e115-24.
65. Marino FE, Cannon J, and Kay D. Neuromuscular responses to hydration in moderate to warm ambient conditions during self-paced high-intensity exercise. *Br J Sports Med*. 2010;44:961-7.
66. Smith MF, Newell AJ, and Baker MR. Effect of acute mild dehydration on cognitive-motor performance in golf. *J Strength Cond Res*. 2012;26:3075-80.
67. Casa DJ, Stearns RL, Lopez RM, Ganio MS, McDermott BP, Walker Yeargin S, Yamamoto LM, Mazerolle SM, Roti MW, Armstrong LE, and Maresh CM. Influence of hydration on physiological function and performance during trail running in the heat. *J Athl Train*. 2010;45:147-56.
68. Skein M and Duffield R. The effects of fluid ingestion on free-paced intermittent-sprint performance and pacing strategies in the heat. *J Sports Sci*. 2010;28:299-307.
69. Hayes LD and Morse CI. The effects of progressive dehydration on strength and power: Is there a dose response? *Eur J Appl Physiol*. 2010;108:701-7.
70. Merry TL, Ainslie PN, and Cotter JD. Effects of aerobic fitness on hypohydration-induced physiological strain and exercise impairment. *Acta Physiol*. 2010;198:179-90.
71. Stearns RL, Casa DJ, Lopez RM, McDermott BP, Ganio MS, Decher NR, Scruggs IC, West AE, Armstrong LE, and Maresh CM. Influence of hydration status on pacing during trail running in the heat. *J Strength Cond Res*. 2009;23:2533-41.
72. Ebert TR, Martin DT, Bullock N, Mujika I, Quod MJ, Farthing LA, Burke LM, and Withers RT. Influence of hydration status on thermoregulation and cycling hill climbing. *Med Sci Sports Exerc*. 2007;39:323-9.
73. Oliver SJ, Laing SJ, Wilson S, Bilzon JL, and Walsh N. Endurance running performance after 48 h of restricted fluid and/or energy intake. *Med Sci Sports Exerc*. 2007;39:316-22.
74. Edwards AM, Mann ME, Marfell-Jones MJ, Rankin DM, Noakes TD, and Shillington DP. Influence of moderate dehydration on soccer performance: Physiological responses to 45 min of outdoor match-play and the immediate subsequent performance of sport-specific and mental concentration tests. *Br J Sports Med*. 2007;41:385-91.
75. Daries HN, Noakes TD, and Dennis SC. Effect of fluid intake volume on 2-h running performances in a 25 degrees c environment. *Med Sci Sports Exerc*. 2000;32:1783-1789.
76. McConell GK, Stephens TJ, and Canny BJ. Fluid ingestion does not influence intense 1-h exercise performance in a mild environment. *Med Sci Sports Exerc*. 1999;31:386-392.
77. Greiwe JS, Staffey KS, Melrose DR, Narve MD, and Knowlton RG. Effects of dehydration on isometric muscular strength and endurance. *Med Sci Sports Exerc*. 1998;30:284-288.
78. Maughan RJ, Fenn CE, and Leiper JB. Effects of fluid, electrolyte and substrate ingestion on endurance capacity. *Eur J Appl Physiol Occup Physiol*. 1989;58:481-6.
79. Kay D and Marino FE. Failure of fluid ingestion to improve self-paced exercise performance in moderate-to-warm humid environments. *J Thermal Biol*. 2003;28:29-34.
80. Dugas JP, Oosthuizen U, Tucker R, and Noakes TD. Rates of fluid ingestion alter pacing but not thermoregulatory responses during prolonged exercise in hot and humid conditions with appropriate convective cooling. *Eur J Appl Physiol*. 2009;105:69-80.
81. Backx K, van Someren KA, and Palmer GS. One hour cycling performance is not affected by ingested fluid volume. *Int J Sport Nutr Exerc Metab* 2003;13:333-342.
82. McLellan TM, Cheung SS, Latzka WA, Sawka MN, Pandolf KB, Millard CE, and Withey WR. Effects of dehydration, hypohydration, and hyperhydration on tolerance during uncompensable heat stress. *Can J Appl Physiol*. 1999;24:349-61.
83. Bachle L, Eckerson J, Albertson L, Ebersole K, Goodwin J, and Petzel D. The effect of fluid replacement on endurance performance. *J Strength Cond Res*. 2001;15:217-24.
84. Sawka MN, Young AJ, Latzka WA, Neuffer PD, Quigley MD, and Pandolf KB. Human tolerance to heat strain during exercise: Influence of hydration. *J Appl Physiol*. 1992;73:368-75.
85. Ali A, Gardiner R, Foskett A, and Gant N. Fluid balance, thermoregulation and sprint and passing skill performance in female soccer players. *Scandinavian Journal of Medicine & Science in Sports*. 2010.
86. Kenefick RW, Cheuvront SN, Palombo LJ, Ely BR, and Sawka MN. Skin temperature modifies the impact of hypohydration on aerobic performance. *J Appl Physiol* (1985). 2010;109:79-86.
87. Sawka MN and Noakes TD. Does dehydration impair exercise performance? *Med Sci Sports Exerc*. 2007;39:1209-1217.
88. Armstrong LE, Johnson EC, and Bergeron MF. Counterview: Is drinking to thirst adequate to appropriately maintain hydration status during prolonged endurance exercise? No. *Wilderness Environ Med*. 2016;27:195-8.
89. Hoffman MD, Cotter JD, Goulet ED, and Laursen PB. View: Is drinking to thirst adequate to appropriately maintain hydration status during prolonged endurance exercise? Yes. *Wilderness Environ Med*. 2016;27:192-5.
90. Goulet ED. Effect of exercise-induced dehydration on time-trial exercise performance: A meta-analysis. *Br J Sports Med*. 2011;45:1149-1156.
91. Goulet ED. Effect of exercise-induced dehydration on endurance performance: Evaluating the impact of exercise protocols on outcomes using a meta-analytic procedure. *Br J Sports Med*. 2013;47:679-86.
92. Savoie FA, Kenefick RW, Ely BR, Cheuvront SN, and Goulet ED. Effect of hypohydration on muscle endurance, strength, anaerobic power and capacity and vertical jumping ability: A meta-analysis. *Sports Med*. 2015;45:1207-27.
93. Goulet ED. Dehydration and endurance performance in competitive athletes. *Nutr Rev*. 2012;70 Suppl 2:S132-6.
94. Barr SI. Effects of dehydration on exercise performance. *Can J Appl Physiol*. 1999;24:164-72.
95. Cheuvront SN, Carter R, III, and Sawka MN. Fluid balance and endurance exercise performance. *Curr Sports Med Rep*. 2003;2:202-208.
96. Cheuvront SN, Kenefick RW, Mountain SJ, and Sawka MN. Mechanisms of aerobic performance impairment with heat stress and dehydration. *J Appl Physiol*. 2010;109:1989-95.
97. Mundel T. To drink or not to drink? Explaining "contradictory findings" in fluid replacement and exercise performance: Evidence from a more valid model for real-life competition. *Br J Sports Med*. 2011;45:2.
98. Beis LY, Wright-Whyte M, Fudge B, Noakes T, and Pitsiladis YP. Drinking behaviors of elite male runners during marathon competition. *Clin J Sport Med*. 2012;22:254-261.
99. Casa DJ, Clarkson PM, and Roberts WO. American college of sports medicine roundtable on hydration and physical activity: Consensus statements. *Curr Sports Med Rep*. 2005;4:115-127.
100. Institute of Medicine. Dietary reference intakes for water, potassium, sodium, chloride, and sulfate. Washington, D.C.: National Academies Press; 2004.
101. Garth AK and Burke LM. What do athletes drink during competitive sporting activities? *Sports Med*. 2013;43:539-64.
102. Rehrer NJ, Brouns F, Beckers EJ, ten HF, and Saris WH. Gastric emptying with repeated drinking during running and bicycling. *Int J Sports Med*. 1990;11:238-243.
103. Vist GE and Maughan RJ. The effect of osmolality and carbohydrate content on the rate of gastric emptying of liquids in man. *J Physiol*. 1995;486:523-531.
104. Gisolfi CV, Summers RW, Lambert GP, and Xia T. Effect of beverage osmolality on intestinal fluid absorption during exercise. *J Appl Physiol*. 1998;85:1941-1948.
105. Costill DL, Kammer WF, and Fisher A. Fluid ingestion during distance running. *Arch Environ Health*. 1970;21:520-525.
106. Speedy DB, Noakes TD, Kimber NE, Rogers IR, Thompson JM, Boswell DR, Ross JJ, Campbell RG, Gallagher PG, and Kuttner JA. Fluid balance during and after an ironman triathlon. *Clin J Sport Med*. 2001;11:44-50.
107. Cosgrove SD, Love TD, Brown RC, Baker DF, Howe AS, and Black KE. Fluid and electrolyte balance during two different preseason training sessions in elite rugby union players. *J Strength Cond Res*. 2014;28:520-7.
108. Kraft JA, Green JM, Bishop PA, Richardson MT, Neggers YH, and Leeper JD. The influence of hydration on anaerobic performance: A review. *ResQExercSport*. 2012;83:282-292.